

Abridged Prescribing Information

Cream with Beclomethasone, Phenylephrine & Lignocaine

Proctosedyl® bd

COMPOSITION:

Beclomethasone Dipropionate IP 0.025% w/w + Phenylephrine Hydrochloride IP 0.1% w/w + Lignocaine Hydrochloride IP 2.5% w/w + Chlorocresol IP (as preservative) 0.1% w/w in aqueous cream base

Pack: 20gm tube with applicator nozzle

THERAPEUTIC INDICATIONS:

Proctosedyl® bd is indicated for pain and bleeding associated with anal fissures and internal or external haemorrhoids; post haemorrhoidectomy pain; minor degrees of proctitis.

DOSAGE AND ADMINISTRATION: Apply a small amount of cream two or three times a day initially, using the applicator if internal administration is required. When inflammation is subsiding, once-daily application is sufficient in most cases. One tube should provide treatment for at least a week

SAFETY-RELATED INFORMATION:

Contraindications: Corticosteroids should not be used in the presence of infection unless effective chemotherapy is also employed. Hypersensitivity to any component of the preparation.

Precautions: Topical administration of corticosteroids to pregnant animals can cause abnormalities of foetal development. The relevance of this finding to human beings has not been established; however topical steroids should not be used extensively in pregnancy, i.e. in large amounts or for prolonged periods.

Adverse Reactions: Prolonged and intensive treatment with active, corticosteroid preparations may cause local atrophic changes in the skin.

For full prescribing information, refer to the company website www.sanofi.in

Source: Anovate PI (Accessed on May 2026)

Updated: July 2026