



Clinical Trials Appendices

R&D Pipeline – New Molecular Entities(*)

Phase 1 (Total:16)		Phase 2 (Total:13)		Phase 3 (Total:8)	Registration (Total:3)
SAR439794 TLR4 agonist Peanut Allergy	SAR228810 Anti-protofibrillar AB mAb Alzheimer's Disease	SAR440340 (**) Anti-IL33 mAb Asthma	ST400 (8) ZFN Gene Editing Technology Beta thalassemia	isatuximab Anti-CD38 mAb 3L Relapsing Refractory MM (ICARIA)	cemiplimab (**) PD-1 inhibitor mAb Advanced CSCC (U.S./EU)
SAR408701 Maytansin-loaded anti-CEACAM5 mAb Solid Tumors	SAR442168 (4) BTK inhibitor Multiple Sclerosis	SAR156597 IL4/IL13 bi-specific mAb Systemic Sclerosis	SAR422459 ABCA4 gene therapy Stargardt Disease	avalglucosidase alfa Neo GAA Pompe Disease	Zynquista ™(**) Oral SGLT-1&2 inhibitor Type 1 Diabetes (U.S./EU)
SAR439459 anti-TGFβ mAb Advanced Solid Tumors	UshStat ® Myosin 7A gene therapy Usher Syndrome 1B	GZ389988 TRKA antagonist Osteoarthritis	SAR425899 GLP-1/GCG dual agonist Obesity/Overweight in T2D	venglustat Oral GCS inhibitor ADPKD(10)	Cablivi ™ Bivalent anti-vWF Nanobody acquired Thrombotic Thrombocytopenic Purpura (EU)
REGN3767 (1) Anti LAG-3 mAb Advanced Cancers	SAR438335 GLP-1/GIP dual agonist Type 2 Diabetes	Combination ferroquine / OZ439 (**) Antimalarial	SAR407899 rho kinase Microvascular Angina	fitusiran siRNA targeting anti-thrombin Hemophilia A and B	
REGN4659 (1) Anti-CTLA-4 mAb Cancer	SAR440181 (9)(**) Myosin activation Dilated Cardiomyopathy	ALX0171 Anti RSV Nanobody Respiratory Syncytial Virus	HIV Viral vector prime & rgp120 boost vaccine	sutimlimab (11) Anti Complement C1s mAb Cold Agglutinin Disease	
REGN4018 (1) Anti MUC16-CD3 bispecific mAb Ovarian Cancer	SAR247799 S1P1 agonist Cardiovascular indication	olipudase alfa rhASM Acid Sphingomyelinase Deficiency(6)	SP0232 (9) mAb(**) Respiratory syncytial virus Monoclonal Antibody	SAR341402 Rapid acting insulin Type 1/2 Diabetes	
SAR439859 SERD Metastatic Breast Cancer	Herpes Simplex Virus Type 2 HSV-2 vaccine	SAR339375 (7) miRNA-21 Alport Syndrome		efpeglenatide (**) Long-acting GLP-1 agonist Type 2 Diabetes	
BIVV001 (2) rFVIII/Fc – vWF – XTEN(3) Hemophilia A	Respiratory syncytial virus Infants Vaccines			mavacamten (12)(**) Myosin inhibitor Obstructive Hypertrophic Cardiomyopathy	

R Registrational Study (other than Phase 3)

O Opt-in rights products for which rights have not been exercised yet

(1) Regeneron product for which Sanofi has opt-in rights

(2) Sanofi Product for which Sobi has opt-in rights

(3) Recombinant Coagulation Factor VIII Fc – von Willebrand Factor – XTEN Fusion protein

(4) Also known as PRN2246

(5) Also known as MYK491

(6) Also known as Niemann Pick type B

(7) Regulus product for which Sanofi has opt-in rights

(8) Developed in collaboration with Sangamo

(9) Also known as MEDI8897

(10) Autosomal Dominant Polycystic Kidney Disease

(11) Also Known as BIVV009

(12) Also known as SAR439152 and MYK461

(*) Data related to all studies published on clinicaltrials.gov

(**) Partnered and/or in collaboration – Sanofi may have limited or shared rights on some of these products

Immuno-inflammation

Oncology

Rare Diseases

Rare Blood Disorders

MS & Neuro

Diabetes

Cardiovascular & metabolism

Vaccines

Additional Indications(*)

Phase 1 (Total:6)	Phase 2 (Total:16)	Phase 3 (Total:20)	Registration (Total:5)
SAR439459 + cemiplimab^(**) Anti-TGFβ mAb + PD-1 inhibitor mAb Advanced Solid Tumors	dupilumab^(**) Anti-IL4Rα mAb Eosinophilic Esophagitis	cemiplimab^(**) PD-1 inhibitor mAb 2L NSCLC	dupilumab^(**) Anti-IL4Rα mAb Asthma 6 - 11 years old
O cemiplimab^(**) + REGN3767⁽¹⁾ PD-1 inhibitor mAb + anti LAG-3 mAb Advanced Cancers	dupilumab^(**) Anti-IL4Rα mAb Grass Immunotherapy	venlustat Oral GCS inhibitor Fabry Disease	dupilumab^(**) Anti-IL4Rα mAb Nasal Polyposis
O cemiplimab^(**) + REGN4659⁽¹⁾ PD-1 inhibitor mAb + Anti-CTLA-4 mAb NSCLC	R sarilumab^(**) Anti-IL6R mAb Polyarticular Juvenile Idiopathic Arthritis	venlustat Oral GCS inhibitor Gaucher Type 3	Dupixent^{®(**)} dupilumab Atopic Dermatitis 12 – 17 years old
O cemiplimab^(**) + REGN4018⁽¹⁾ PD-1 inhibitor mAb + Anti-MUC16-CD3 bispecific mAb - Ovarian Cancer	sarilumab^(**) Anti-IL6R mAb Systemic Juvenile Arthritis	venlustat Oral GCS inhibitor Gaucher related Parkinson's Disease	Dupixent^{®(**)} dupilumab Atopic Dermatitis 6 – 11 years old
SAR439859 SERD + Palbociclib Metastatic Breast Cancer	SAR440340^(**) Anti-IL33 mAb COPD	mavacamten^{(3)(**)} Myosin inhibitor Non-Obstructive Hypertrophic Cardiomyopathy	Dupixent^{®(**)} dupilumab Atopic Dermatitis 6 months - 5 years old
sutimlimab⁽²⁾ Anti Complement C1s mAb Immune Thrombocytopenia	R cemiplimab^(**) PD-1 inhibitor mAb Advanced Basal Cell Carcinoma	Rabies VRVg Purified vero rabies vaccine	Dupixent^{®(**)} dupilumab Atopic Dermatitis 6 months - 5 years old
	isatuximab + cemiplimab^(**) Anti-CD38 mAb + PD-1 inhibitor mAb Relapsing Refractory MM	Adacel+ Tdap booster	cemiplimab^(**) PD-1 inhibitor mAb + ipilimumab 1L NSCLC
	isatuximab + cemiplimab^(**) Anti-CD38 mAb + PD-1 inhibitor mAb Advanced Malignancies	Shan 6 DTP-HepB-Polio-Hib Pediatric hexavalent vaccine	cemiplimab^(**) PD-1 inhibitor mAb + platinum based chemotherapy 1L NSCLC
		cemiplimab^(**) PD-1 inhibitor mAb 2L Cervical Cancer	cemiplimab^(**) PD-1 inhibitor mAb + platinum based chemotherapy 1L NSCLC
		isatuximab Anti-CD38 mAb 1L Newly Diagnosed MM (IMROZ)	isatuximab Anti-CD38 mAb 1L Newly Diagnosed MM (IMROZ)
			isatuximab Anti-CD38 mAb 1-3L Relapsing Refractory MM (IKEMA)
			Aubagio[®] teriflunomide Relapsing Multiple Sclerosis - Pediatric
			Lemtrada[®] alemtuzumab Relapsing Remitting Multiple Sclerosis - Pediatric
			Zynquista^{TM(**)} Oral SGLT-1&2 inhibitor Type 2 Diabetes
			Zynquista^{TM(**)} Oral SGLT-1&2 inhibitor Worsening Heart Failure in Diabetes
			Cerdelga[®] eliglustat Gaucher Type 1, switch from ERT - Pediatric
			Praluent^{®(**)} alirocumab LDL-C reduction - Pediatric
			Fluzone[®] QIV HD Quadrivalent inactivated Influenza vaccine - High dose
			Men Quad TT Advanced generation meningococcal ACYW conjugate vaccine
			Pediatric pentavalent vaccine DTP-Polio-Hib Japan
			dupilumab^(**) Anti-IL4Rα mAb Asthma 12y+ (U.S./EU)
			Praluent^{®(**)} alirocumab CV events reduction (U.S./EU)
			VaxiGrip[®] QIV IM Quadrivalent inactivated Influenza vaccine 6 - 35 months
			PR5i DTP-HepB-Polio-Hib Pediatric hexavalent vaccines (U.S.)
			Fluzone[®] 0.5 mL QIV Quadrivalent inactivated Influenza vaccine 6 months+

R Registrational study (other than Phase 3)

O Opt-in rights products for which rights have not been exercised yet

- (1) Regeneron product for which Sanofi has opt-in rights
 (2) Also known as BIVV009
 (3) Also known as SAR439152 and MYK461
 (4) U.S. filing pending acceptance by FDA
 (*) Data related to all studies published on clinicaltrials.gov
 (**) Partnered and/or in collaboration - Sanofi may have limited or shared rights on some of these products

Expected Submission Timeline⁽¹⁾

NMEs

isatuximab anti-CD38 mAb 3L RRRM (ICARIA)	SAR341402 Rapid acting insulin Type 1/2 Diabetes - EU ⁽²⁾	avalglucosidase alfa Neo GAA Pompe Disease	olipudase alfa rhASM ASD ⁽⁴⁾	fitusiran siRNA inhibitor Hemophilia A/B	sutimlimab⁽⁵⁾ Anti Complement C1s mAb Cold Agglutinin Disease	venglustat Oral GCS inhibitor ADPKD ⁽⁶⁾	efpeglenatide⁽⁷⁾ Long acting GLP1-R agonist Type 2 Diabetes	GZ389988 TRKA antagonist Osteoarthritis	ALX0171 Anti RSV Nanobody Respiratory Syncytial Virus	SAR156597 IL4/IL13 bi-specific mAb Systemic Scleroderma	SAR228810 Anti-protofibrillar AB mAb Alzheimer's Disease	SAR407899 rho kinase Microvascular Angina	SAR440340⁽⁸⁾ Anti-IL33 mAb Asthma	SAR422459 ABCA4 gene therapy Stargardt Disease	SP0232 mAbs⁽⁷⁾⁽⁸⁾ Respiratory syncytial virus	SAR425899 GLP-1/GCG dual agonist Obesity/Overweight in T2D	HIV Viral vector prime & rgp120 boost vaccine
--	---	---	--	---	---	---	---	--	--	--	---	--	---	---	--	---	--

Additional Indications

2018	2019	2020	2021	2022 and beyond
Dupixent[®](**) dupilumab AD 12 – 17 years old	Dupixent[®](**) dupilumab AD 6 - 11 years old -	Zynquista[™](**) Oral SGLT-1&2 inhibitor Type 2 Diabetes – EU ⁽³⁾	sarilumab^(**) Anti-IL6R mAb Polyarticular Juvenile Idiopathic Arthritis	Aubagio[®] teriflunomide Relapsing MS – Ped.
	dupilumab^(**) Anti-IL4Rα mAb Nasal Polyposis Adult	Fluzone[®] QIV HD Quadrivalent inactivated Influenza vaccine - High dose	cemiplimab^(**) PD-1 inhibitor mAb 2L Cervical Cancer	Shan 6 DTP-HepB-Polio-Hib Pediatric hexavalent vaccine
	cemiplimab^(**) PD-1 inhibitor mAb Advanced BCC	Men Quad TT Adv. generation meningococcal U.S.: 2y+ & EU: Toddlers+	isatuximab Anti-CD38 mAb 1-3L RRRM (IKEMA)	isatuximab Anti-CD38 mAb (IMROZ) 1L Newly Diagnosed MM
	Pentacel[®] vIPV DTaP-IPV/Hib		cemiplimab^(**) PD-1 inhibitor mAb 1L NSCLC	cemiplimab^(**) PD-1 inhibitor mAb + platinum based chemotherapy 1L NSCLC
			Adacel+ Tdap booster	Zynquista[™](**) Oral SGLT-1&2 inhibitor Worsening Heart Failure in Diabetes
			Pediatric pentavalent vaccine DTP-Polio-Hib (Japan)	Adacel+ Tdap booster
				Dupixent[®](**) dupilumab AD 6 months - 5 years old
				sarilumab^(**) Anti-IL6R mAb Systemic Juvenile Arthritis
				SAR440340^(**) Anti-IL33 mAb COPD
				venglustat Oral GCS inhibitor Fabry Disease
				Praluent[®](**) alirocumab LDL-C reduction - Pediatric
				dupilumab^(**) Anti-IL4Rα mAb Eosinophilic Esophagitis
				venglustat Oral GCS inhibitor Gaucher Type 3
				dupilumab^(**) Anti-IL4Rα mAb Asthma 6 - 11 years old
				Rabies VRVg Purified vero rabies vaccine
				Cerdelga[®] eliglustat Gaucher Type 1, switch from ERT – Pediatric EU

 Immuno-inflammation	 MS & Neuro
 Oncology	 Diabetes
 Rare Diseases	 Cardiovascular & metabolism
 Rare Blood Disorders	 Vaccines

(1) Excluding Phase 1 - Data related to all studies published on clinicaltrials.gov
 (2) Submission strategy for the U.S. under evaluation
 (3) Submission for the U.S. expected in 2020
 (4) Acid Sphingomyelinase Deficiency
 (5) Also known as BIVV009; Currently operating as separate entities. Reported dates are based on prior Bioverativ disclosure of study completion date
 (6) Autosomal Dominant Polycystic Kidney Disease
 (7) Also known as MEDI8897
 (8) Gaucher Related Parkinson's Disease
 (***) Partnered and/or in collaboration – Sanofi may have limited or shared rights on some of these products

Pipeline Movements Since Q1 2018

	Additions	Removals	
Registration	<p>Cabliivi™ Bivalent anti-vWF Nanobody acquired Thrombotic Thrombocytopenic Purpura (EU)</p> <p>Praluent®^(**) alirocumab CV events reduction (U.S.⁽¹⁾/EU)</p>		
Phase 3	<p>mavacamten^{(2)(**)} Myosin inhibitor Obstructive Hypertrophic Cardiomyopathy</p>	<p>Zynquista™^(**) SGLT 1 & 2 inhibitor Worsening Heart Failure in Diabetes</p>	
	<p>Praluent®^(**) alirocumab LDL-C reduction - Pediatric</p>	<p>cemiplimab^(**) PD-1 inhibitor mAb + ipilimumab 1L NSCLC</p>	
	<p>Cerdelga® eliglustat Gaucher disease Type 1, switch from ERT Pediatric</p>	<p>cemiplimab^(**) PD-1 inhibitor mAb + platinum based chemotherapy 1L NSCLC</p>	
	<p>venglustat Oral GCS inhibitor ADPKD⁽³⁾</p>		
Phase 2	<p>dupilumab^(**) Anti-IL4Ra mAb Grass Immunotherapy</p>	<p>ST400⁽⁴⁾ ZFN Gene Editing Technology Beta thalassemia</p>	<p>SAR566658 Maytansin-loaded anti-CA6 mAb Triple Negative Breast Cancer</p>
	<p>SAR440340^(**) Anti-IL33 mAb COPD</p>	<p>mavacamten^{(2)(**)} Myosin inhibitor Non-Obstructive Hypertrophic Cardiomyopathy</p>	<p>Tuberculosis Recombinant subunit vaccine</p>
	<p>ALX0171 Anti RSV nanobody Respiratory Syncytial Virus</p>	<p>cemiplimab^(**) PD-1 inhibitor mAb 2L NSCLC</p>	
Phase 1	<p>REGN4659⁽⁵⁾ Anti-CTLA-4 mAb Cancer</p>	<p>cemiplimab^(**) + REGN4659⁽⁵⁾ PD-1 inhibitor mAb + Anti-CTLA-4 mAb NSCLC</p>	
	<p>REGN4018⁽⁵⁾ Anti MUC16-CD3 bispecific mAb Ovarian Cancer</p>	<p>cemiplimab^(**) + REGN4018⁽⁵⁾ PD-1 inhibitor mAb + Anti MUC16-CD3 bispecific mAb - Ovarian Cancer</p>	

R&D Pipeline Summary – Total Projects⁽¹⁾

	Phase 1	Phase 2	Phase 3	Registration	TOTAL
Immuno-inflammation	1	10	5	1	17
Oncology	11	4	7	1	23
Rare Diseases	0	4	3	0	7
Rare Blood Disorders	2	1	2	1	6
Multiple Sclerosis and Neurology	3	2	2	0	7
Diabetes	1	1	4	1	7
Cardiovascular Diseases	2	2	2	1	7
Vaccines	2	5	3	3	13
TOTAL	22	29	28	8	

51 **36**

87 Total Projects

Expected R&D Milestones

Products	Expected milestones	Timing
Praluent®	U.S. sBLA filing to include ODYSSEY OUTCOMES results ⁽¹⁾	Q3 2018
isatuximab	Start of Phase 3 in 1 st line Multiple Myeloma in SCT eligible patients (GMMG)	Q3 2018
Cablivi™ (caplacizumab)	U.S. FDA filing in acquired Thrombotic Thrombocytopenic Purpura	Q3 2018
venglustat	Start of Pivotal study in Autosomal Dominant Polycystic Kidney Disease	Q3 2018
MenQuadTT	Phase 3 results for prevention of Meningococcal Meningitis	Q3 2018
Dupixent®	U.S. FDA filing in Atopic Dermatitis in Adolescent patients	Q3 2018
Fluzone® QIV HD	Phase 3 results for prevention of Influenza	Q4 2018
cemiplimab	U.S. regulatory decision in locally advanced CSCC	Q4 2018
dupilumab	U.S. regulatory decision in Asthma in Adult/Adolescent patients	Q4 2018
dupilumab	Start of Phase 2b/3 trial in Chronic Obstructive Pulmonary Disease	Q4 2018
isatuximab	Phase 3 results in Multiple Myeloma in combination with PomDex (ICARIA)	Q4 2018
dupilumab	Phase 3 read-out in Nasal Polyps	Q4 2018
dupilumab	Start of Phase 3 trial in Eosinophilic Esophagitis	Q4 2018
efpeglenatide	Start of Phase 3 in Type 2 Diabetes as add-on to metformin vs dulaglutide	Q4 2018
efpeglenatide	Start of Phase 3 in Type 2 Diabetes as add-on to basal insulins	Q4 2018
alemtuzumab	Start of Phase 3 in Primary Progressive Multiple Sclerosis	H2 2018
Zynquista™ (sotagliflozin)	EU CHMP decision expected in Type 1 Diabetes	Q1 2019
cemiplimab	EU CHMP decision expected in Advanced Cutaneous Squamous Cell Carcinoma	Q1 2019

List of abbreviations

AE	Adverse Events	IAR	Infusion Associated Reaction	QNM	Every N Months
APO	Apolipoprotein	IC	Investigator's Choice	QOL	Quality Of Life
BOR	Best Overall Response	IGA	Investigator's Global Assessment	RECIST	Response Evaluation Criteria in Solid Tumors
BW	Body Weight	IMiD	Immunomodulatory Drug	SAE	Serious Adverse Events
CB	Clinical Benefit	ITT	Intent To Treat	SBP	Systolic Blood Pressure
CNS	Central Nervous System	LP	Lipoprotein	SDMT	Symbol Digit Modalities Test
CR	Complete Response	MRI	Magnetic Resonance Imaging	SMPG	Self Monitored Plasma Glucose
CRR	Complete Response Rate	MTD	Maximum Tolerated Dose	SSD	Study Start Date
CT	Computed Tomography	N	Number	TC	Total Cholesterol
CV	Cardiovascular	NC	Nasal Congestion/obstruction	TEAE	Treatment Emergent Adverse Events
DE	Data Expected	NNT	Number Needed to Treat	TSS	Total Symptom Score
DCR	Disease Control Rate	OS	Overall Survival	TG	Triglycerides
DLT	Dose-Limiting Toxicity	ORR	Overall Response Rate	TTP	Time To Progression
DOD	Duration Of Disease	PD	Pharmacodynamics	TTR	Time To Response
DOR	Duration Of Response	PI	Proteasome Inhibitor	TX	Treatment
DPP4	Dipeptidyl peptidase 4	PFS	Progression-Free Survival	VGPR	Very Good Partial Response
EASI	Eczema Area and Severity Index	PK	Pharmacokinetic		
FPG	Fasting Plasma Glucose	PPG	Postprandial Glucose		
HbA1c	Hemoglobin A1c	PRO	Patient Reported Outcome		
IAE	Incidence of Adverse Events	QNW	Every N Weeks		

Dupilumab (anti-IL4R α mAb) Asthma 1/2

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
LIBERTY ASTHMA TRAVERSE LTS12551 NCT02134028	Phase 2/3 Open label extension study long-term safety & tolerability evaluation in patients with asthma who participated in previous studies	2,284 enrolled	<ul style="list-style-type: none"> For patients coming from DRI12544, PDY14192, EFC13579, EFC13691 studies, added to current controller medications Open-label, 	<ul style="list-style-type: none"> Primary: N and % of patients experiencing any TEAE Secondary: Safety 	<ul style="list-style-type: none"> SSD: Jul. 2014 DE: 2019
LIBERTY ASTHMA EXCURSION LTS14424 NCT03560466	Phase 3 Long term safety and tolerability (1 year) of dupilumab in pediatric patients with asthma who participated in a previous dupilumab asthma clinical study	377	<ul style="list-style-type: none"> Open-label 1 year of Tx 	<ul style="list-style-type: none"> Primary: N of patients experiencing any TEAE Secondary: Severe asthma exacerbation events, change in % predicted FEV1, in absolute FEV1, in FVC, FEF, dupilumab concentrations, anti-dupilumab Ab, eosinophils, Ig, IgE 	<ul style="list-style-type: none"> SSD: June 2018 DE: 2026

Dupilumab (anti-IL4R α mAb) Asthma 2/2

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
CHILDREN ASTHMA VOYAGE EFC14153 NCT02948959	Phase 3 Evaluation of dupilumab in children (6 to <12 years) with uncontrolled asthma	471	<ul style="list-style-type: none"> In children 6 to <12 years of age with uncontrolled persistent asthma Randomized, Double-blind, Placebo-controlled, parallel group 52 weeks Tx, 12 weeks post Tx 	<ul style="list-style-type: none"> Primary: Annualized rate of severe exacerbation events during Tx period Secondary: Safety and tolerability, PROs, Systemic exposure and incidence of anti-drug antibodies, Association between dupilumab Tx and pediatric immune responses to vaccines 	<ul style="list-style-type: none"> SSD: Jun. 2017 DE: 2021

Dupilumab (anti-IL4R α mAb) Atopic Dermatitis (AD) 1/3

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
OLE Pediatrics AD R668-AD-Reg 1434 NCT02612454	Phase 3 A study to assess the long-term safety of dupilumab administered in patients 6 to <18 years of age with AD	765 expected	<ul style="list-style-type: none"> For patients having participated in a prior dupilumab study in pediatrics with AD Open label extension study 	<ul style="list-style-type: none"> Primary: Incidence and rate of TEAEs Secondary: SAEs and TEAEs of special interest, % of patients who achieve and maintain remission, EASI-75: % of patients achieving and maintaining at least 75% reduction in EASI score over time, EASI-50: % of patients achieving and maintaining at least 50% reduction in EASI scores over time 	<ul style="list-style-type: none"> SSD: Oct. 2015 DE: 2023
Pediatrics (12 to 17 years) AD R668-AD-Reg 1526 NCT03054428	Phase 3 A study to investigate the efficacy and safety of dupilumab monotherapy in patients 12 to 17 years of age, with moderate-to-severe AD	251	<ul style="list-style-type: none"> Pediatric patients (12 to 17 years old) with moderate-to-severe AD A randomized, double-blind, placebo-controlled, 3-arm: dupilumab dose 1, dupilumab dose 2, placebo 	<ul style="list-style-type: none"> Primary: % of patients with IGA 0 to 1 (on a 5-point scale), % of patients with EASI-75 Secondary: % change in EASI score 	<ul style="list-style-type: none"> SSD: Apr. 2017 DE: 2018

Dupilumab (anti-IL4R α mAb) Atopic Dermatitis (AD) 2/3

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
LIBERTY AD PRESCHOOL R668-AD-1539 NTC03346434	Phase 2/3 Safety, Pharmacokinetics and Efficacy of Dupilumab in Patients \geq 6 Months to <6 Years With Severe Atopic Dermatitis	280	<ul style="list-style-type: none"> Part A: Open-label, single-ascending dose, sequential cohort phase 2 study Part B: Randomized, double-blind, parallel-group, placebo-controlled phase 3 study 	<ul style="list-style-type: none"> Part A: PK Part B: Proportion of patients with Investigator's Global Assessment "0" or "1" (on a 5-point scale) at week 16 	<ul style="list-style-type: none"> SSD: Dec. 2017 DE: 2022
AD in 6 - 11 Years Old R668-AD-1652 NCT03345914	Phase 3 Efficacy and safety of Dupilumab administered with Topical Corticosteroids in participants \geq 6 to <12 years with Severe Atopic Dermatitis	240	<ul style="list-style-type: none"> Randomized, Double-blind, Placebo-controlled Study 	<ul style="list-style-type: none"> Primary: Proportion of patients with Investigator's Global Assessment "0" or "1" (on a 5-point scale) at week 16 Secondary: Change from baseline to week 16 in Children's Dermatology Life Quality Index, Percent change in EASI score from baseline to week 16, Incidence of serious TEAEs through week 16 	<ul style="list-style-type: none"> SSD: Jan. 2018 DE: 2019

Dupilumab (anti-IL4R α mAb) Atopic Dermatitis (AD) 3/3

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
Autoinjector R668-AD-1607 NTC03050151	Phase 1 Study of Dupilumab Auto-injector Device When Used by Patients With Atopic Dermatitis	176	<ul style="list-style-type: none"> Part A: Patients with moderate-to-severe AD will be randomized to receive dupilumab (dose 1) by auto-injector (AI) device or prefilled syringe. Part B: Once part A is completely enrolled, part B will randomize patients with moderate-to-severe AD to receive dupilumab (dose 2) by auto-injector (AI) device or prefilled syringe. 	<ul style="list-style-type: none"> Primary: Number and type of validated AI device-associated PTFs during the treatment period by actual number of injections Secondary: Number of patients with an AI device associated PTF, Number and type of AI device-associated PTCs, Number of patients with an AI device associated PTC, Type of AI device-associated failed drug deliveries, Number of patients with an AI device-associated failure to deliver dose, PK 	<ul style="list-style-type: none"> SSD: Mar. 2017 DE: 2018
Open-Label R668-AD-1225 NCT01949311	Phase 3 Open-Label study of Dupilumab in patients with Atopic Dermatitis	2733	<ul style="list-style-type: none"> Open label extension study for patients who participated in placebo-controlled dupilumab AD trials. The study primarily evaluates long term safety (adverse events) and immunogenicity. Efficacy parameters are based on IGA, EASI) and the NRS. 	<ul style="list-style-type: none"> Primary: TEAEs, Secondary: SAEs and AEs of special interest, % of patients who achieve and maintain remission, EASI-75: % of patients achieving and maintaining at least 75% reduction in EASI score over time, EASI-50: % of patients achieving and maintaining at least 50% reduction in EASI scores over time 	<ul style="list-style-type: none"> SSD: Oct. 2013 DE: 2018

Dupilumab (anti-IL4R α mAb) Nasal Polyposis (NP)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
NP SINUS-24 EFC14146 NCT02912468	Phase 3 Evaluation of dupilumab in patients with bilateral NP on a background of mometasone furoate nasal spray	276	<ul style="list-style-type: none"> Patients with bilateral sinonasal polyposis that despite prior Tx with systemic corticosteroids have an endoscopic bilateral NPS with a score at least of 5 over 8 Randomized, double-blind, placebo-controlled study, 4 weeks run-in, 24 weeks Tx, 24 weeks post-Tx 	<ul style="list-style-type: none"> Primary: NC symptom severity score based on the patient daily morning assessment & by endoscopy Secondary: TSS, Loss of smell, Sinus opacification 	<ul style="list-style-type: none"> SSD: Dec. 2016 DE: 2018
LIBERTY NP SINUS-52 EFC14280 NCT02898454	Phase 3 Evaluation of dupilumab in patients with bilateral NP on a background of mometasone furoate nasal spray	448	<ul style="list-style-type: none"> Patients with bilateral sinonasal polyposis that despite prior Tx with systemic corticosteroids have an endoscopic bilateral NPS with a score at least of 5 over 8 Randomized, double-blind, placebo-controlled study, 4 weeks run-in, 52 weeks Tx, 12 weeks post-Tx, 3-arm, dupilumab dose regimen 1, dupilumab dose regimen 2, placebo 	<ul style="list-style-type: none"> Primary: NC symptom severity score based on the patient daily morning assessment & by endoscopy, Secondary: TSS, Loss of smell, Sinus opacification 	<ul style="list-style-type: none"> SSD: Dec. 2016 DE: 2018

Dupilumab (anti-IL4R α mAb) Grass Immunotherapy

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
GRASS R668 – ALG - 16115 NCT03558997	Phase 2 Evaluation of dupilumab as an adjunct for subcutaneous grass immunotherapy to reduce provoked allergic rhinitis symptoms	100	<ul style="list-style-type: none"> Patients with history of grass pollen-induced seasonal allergic rhinitis confirmed by SPT with Timothy grass extract and Timothy grass specific IgE, Randomized, double-blind, placebo-controlled study, 4 arms: dupilumab + Timothy Grass SCIT; placebo dupilumab + SCIT; dupilumab + placebo SCIT; placebo dupilumab + placebo SCIT; 16 weeks of Tx 	<ul style="list-style-type: none"> Primary: Total Nasal Symptom Score (TNSS) after nasal allergen challenge (NAC) with Timothy grass extract, Secondary: change from baseline in TNSS AUC post NAC over the 1st hour after the challenge across the arms, specific immunoglobulin G4, TEAEs 	<ul style="list-style-type: none"> SSD: June 2018 DE: 2019

Sarilumab (anti-IL6 mAb) Rheumatoid Arthritis (RA)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
SARIL-RA-EXTEND LTS11210 NCT01146652	Phase 3 Long-term evaluation of sarilumab in RA patients	2000	<ul style="list-style-type: none"> In patients with RA having participated to previous trials Multi-center, uncontrolled extension, open-label; up to 1 week screening, at least 264 weeks of Tx to 516 weeks max., 6 weeks post-Tx 	<ul style="list-style-type: none"> Primary: N of patients with AE Secondary: Long term efficacy of sarilumab in patients with RA (ACR20, DAS28, EULAR response) 	<ul style="list-style-type: none"> SSD: Jun. 2010 DE: 2020

Sarilumab (anti-IL6 mAb) Juvenile Idiopathic Arthritis (JIA)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
Polyarticular JIA Children & Adolescents DRI13925 NCT02776735	Phase 2b Dose-finding study of sarilumab in children and adolescents with Polyarticular-course Juvenile Idiopathic Arthritis (pcJIA)	72	<ul style="list-style-type: none"> In children and adolescents, Aged 2 to 17 years, with pcJIA Open-label, sequential, ascending, repeated dose-finding Study; 4-week screening, 12-week core Tx, 144-week extension, 6-week post-Tx 	<ul style="list-style-type: none"> Primary: PK parameters (Up to week 12) Secondary: PD profile, The efficacy and the safety of sarilumab in patients with pcJIA, Long-term safety of sarilumab in patients with pcJIA 	<ul style="list-style-type: none"> SSD: Sep. 2016 DE: Primary completion 2018; full completion 2022
Systemic JIA Children & Adolescents DRI13926 NCT02991469 ⁽¹⁾	Phase 2b Dose-finding study of sarilumab in children and adolescents with Systemic Juvenile Idiopathic Arthritis (sJIA)	36	<ul style="list-style-type: none"> In children and adolescents, aged 1 to 17 years, with sJIA Open-label, sequential, ascending, repeated dose finding study, 4-week screening, 12-week Tx, 92- week extension, 6-week post-Tx 	<ul style="list-style-type: none"> Primary: PK parameters (Up to week 12) Secondary: PD profile, efficacy and the safety of sarilumab in patients with sJIA, Long term safety of sarilumab in patients with sJIA 	<ul style="list-style-type: none"> SSD: Dec. 2017 DE ^(1st part) (2): 2018

(1) Recruitment currently suspended to optimize study design and procedures
 (2) Final Data Collection date for primary outcome measure

SAR156597 (anti-IL13/IL4 mAb) Scleroderma

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
POC in Scleroderma ACT14604 NCT02921971	Phase 2a Efficacy and safety of SAR156597 in the Tx of Diffuse Cutaneous Systemic Sclerosis (dcSSc)	94	<ul style="list-style-type: none"> Randomized, double-blind, Parallel Assignment, placebo-controlled, 4-week screening, 24-week Tx period, 11-week follow-up 	<ul style="list-style-type: none"> Primary: Change from baseline in mRSS Secondary: Change from baseline in Health Assessment Questionnaire Disability Index (HAQ-DI), assessed with SHAQ, Change from baseline in respiratory function as measured by observed Forced Vital Capacity Change from baseline in observed Carbon Monoxide Diffusing Lung Capacity (DLco [corrected for hemoglobin]) 	<ul style="list-style-type: none"> SSD: Nov. 2016 DE (1st part)⁽¹⁾: 2018; full completion 2019

SAR440340 (Anti-IL33 mAb)

Asthma – single agent and in combination with dupilumab

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
Asthma R3500-AS-1619 NCT02999711	Phase 1 Safety and tolerability of multiple ascending subcutaneous doses of SAR440340 in adult patients with Moderate Asthma	23	<ul style="list-style-type: none"> Randomized, double-blind, Placebo-controlled, Multiple ascending dose study of the safety Cohort 1: SAR440340 low dose or placebo Cohort 2 : SAR440340 medium dose or placebo 	<ul style="list-style-type: none"> Primary: Incidence of TEAEs after repeat subcutaneous administration, severity of TEAEs Secondary: Concentration-time profile of REGN3500 after repeat subcutaneous administration, Immunogenicity, % change in total from baseline forced expiratory volume 	<ul style="list-style-type: none"> SSD: Feb. 2017 DE: 2018
Asthma in combination with dupilumab R3500-AS-1633 NCT03112577	Phase 1 Effects of SAR440340 dupilumab, combination of both on markers of inflammation after bronchial allergen challenge in patients with Allergic Asthma	38	<ul style="list-style-type: none"> Patients with mild allergic asthma for at least 6 months, Randomized, Placebo –controlled, Parallel Assignment 5 arms: SAR440340 alone, dupilumab alone, SAR440340 + dupilumab, placebo and fluticasone propionate (active comparator, open label dosing) 	<ul style="list-style-type: none"> Primary: Difference in bronchial allergen challenge (BAC)-induced changes in sputum inflammatory markers in individuals treated with SAR440340, dupilumab and the combination of both, or placebo [Screening (pre-treatment) to week 4 after treatment initiation] Secondary: TEAEs (incidence and severity), PK profile, immunogenicity, difference in the BAC-induced changes in sputum inflammatory mRNA signature in individual patients treated with fluticasone 	<ul style="list-style-type: none"> SSD: July 2017 DE: 2020 (completion)

SAR440340 (Anti-IL33 mAb) Asthma Combination with dupilumab

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
Asthma SA and combination with dupilumab ACT15102 NCT03387852	Phase 2 Efficacy, Safety and Tolerability (POC) of SAR440340 and the coadministration with dupilumab in patients with Moderate-to-severe Asthma, Not Well Controlled on Inhaled Corticosteroid (ICS) Plus Long-acting β 2 Adrenergic Agonist (LABA) Therapy	240	<ul style="list-style-type: none"> Adults patient with a physician diagnosis of asthma for at least 12 months, Randomized, Double-blind, Placebo-controlled, Parallel Group, with fluticasone w/wo salmeterol Arm 1: SAR440340 monotherapy Arm 2 : dupilumab monotherapy Arm 3: coadministration of SAR440340 and dupilumab Arm 4: placebo Ttmt every 2 weeks for 12 weeks Total duration for one patient: appr. 36 weeks, including 4 weeks screening, 12 weeks ttmt and 20 weeks post-ttmt 	<ul style="list-style-type: none"> Primary: LOAC (lost of asthma control) events Secondary: change in FEV1 (forced expiratory volume 1) 	<ul style="list-style-type: none"> SSD: Mar. 2018 DE: 2019

SAR440340 (Anti-IL33 mAb) COPD

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
POC in COPD ACT15104 NCT03546907	Phase 2 Efficacy, Safety and Tolerability (POC) of SAR440340 in patients with moderate-to-severe COPD	340	<ul style="list-style-type: none"> Adults patients with a diagnosis of moderate-to-severe COPD for at least 1 year Randomized, Double-blind, Placebo-controlled, on top of standards of care Arm 1: SAR440340 Arm 2 : placebo Total duration for one patient: 46 to 76 weeks including 10 days to 4 weeks of screening, 24 to 52 weeks Tx period and 20 weeks post IMP Tx period 	<ul style="list-style-type: none"> Primary: AECOPD (Acute Exacerbations in COPD) Secondary: change in pre-bronchodilator FEV1 (forced expiratory volume 1), time to 1st COPD exacerbations, AEs, change in post-bronchodilator FEV1 	<ul style="list-style-type: none"> SSD: Jul. 2018 DE: 2020

SAR439794 (TLR4 agonist) Immunomodulator

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
Peanut Allergy TDR14287 NCT03463135	Phase 1 Safety, Tolerability and Pharmacodynamics of SAR439794 in Peanut Allergic Adult Patients	44	<ul style="list-style-type: none"> Randomized, Double-blind, Placebo-controlled, 3 Arms Repeated Sublingual daily Administration of SAR439794 or placebo Total study duration per participant: approximately from 15 to 18 weeks (core study) from screening until end-of-study visit, and 2 phone calls at Week 26 and Week 52 after the last IMP dose. 	<ul style="list-style-type: none"> Primary: Incidence of AEs Secondary: PD parameters (peanut-specific serum IgG levels, peanut-specific serum IgE levels, SkinPrick test) 	<ul style="list-style-type: none"> SSD: May 2018 DE: 2020 (completion)

Ferroquine – Artefenomel / OZ439

Malaria

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
FALCI DRI12805 NCT02497612	Phase 2 Efficacy, Safety, Tolerability and PK of a single dose regimen of ferroquine with artefenomel (OZ439) in adults and children with Uncomplicated Plasmodium Falciparum Malaria	662	<ul style="list-style-type: none"> Patients from 6 months to 70 years suffering from mono-infection by P. falciparum Randomized, Double-blind, Parallel Assignment 4 doses of ferroquine associated to 1 dose of artefenomel according to age and body weight Study duration: up to 67 days for each patient 	<ul style="list-style-type: none"> Primary: % of patients with Polymerase Reaction Chain (PCR)-adjusted Adequate Clinical and Parasitological Response (ACPR) Secondary: Time to re-emergence, Time to recrudescence, Parasite clearance time, % of patients with PCR - crude ACPR, SAE, AESI, TEAE, % of patients with PCR - adjusted ACPR 	<ul style="list-style-type: none"> SSD: Jul. 2015 DE: 2019

ALX0171

RSV lower respiratory tract infection (LRTI)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
ALX0171-C204 NCT03468829	Phase 2 Efficacy and safety of ALX-0171 in adults with respiratory syncytial virus (RSV) respiratory tract infection who have undergone hematopoietic stem cell transplantation (HSCT). Clinical activity, PK, virology, and immunogenicity of inhaled ALX 0171 also to be assessed	75	<ul style="list-style-type: none">Adults, 18 to 75 years, diagnosed with RSV tract infection after HSCTRandomized, Double-Blind, Placebo-Controlled, Parallel Assignment, 3 arms: 2 doses of ALX-0171 and placeboStudy duration: 42 days; Tx period: oral inhalation once daily for a max. of 14 days, follow up period of 28 days	<ul style="list-style-type: none">Primary: Time-weight avg. change in RSV nasal viral loadSecondary: Incidence of AE, nasal RSV load parameter, clinical stabilization, # of days with/without oxygen supplementation, progression to lower resp. tract disease (LRT) in subjects with upper resp. tract infection (URTI), serum concentration of ALX-0171, and immunogenicity	<ul style="list-style-type: none">SSD: Jun. 2018DE: 2020

ALX0171

RSV LRTI

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
<p>Japan</p> <p>ALX0171-C203 NCT03418571</p>	<p>Phase 2</p> <p>Evaluate the safety, tolerability and systemic PK of ALX0171 in Japanese infants and young children with respiratory syncytial virus (RSV) lower respiratory tract infection (LRTI). Antiviral effect, clinical activity, PD and immunogenicity of inhaled ALX 0171 also to be assessed</p>	60	<ul style="list-style-type: none"> • Infants and young children up to 2 years, hospitalized with RSV LRTI • Randomized, Double-Blind, Placebo-Controlled, Sequential Assignment, 5 arms: 4 doses of ALX0171 and placebo • Study duration: 28 days; Tx period: oral inhalation once daily for 3 days, follow up period of 28 days after 1st dose 	<ul style="list-style-type: none"> • Primary: safety and tolerability measured by serious AE • Secondary: PK, time for viral load to drop below assay quantification limit (BQL), Global Severity Score (sum of 7 scores from feeding, medical interventions, respiratory distress, respiratory rate, apnea, general appearance and body temp), time to clinical response, and immunogenicity 	<ul style="list-style-type: none"> • SSD: Mar. 2018 • DE¹: 2019

Isatuximab (anti-CD38 mAb) Hematological Malignancies (HM)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
CD38+HM TED10893 NCT01084252	Phase1/2 Dose escalation and efficacy study of isatuximab in patients with selected CD38+ HM	346	<ul style="list-style-type: none"> Phase 1: MTD Phase 2: Stage 1: isatuximab activity at different doses/schedules and to select dose and regimen as single agent or in combination with dexamethasone Stage 2: activity at the selected dose/schedule from stage1, as single agent (ISA arm) and in combination with dexamethasone (ISAdex arm) Randomized, Open-label, Parallel assignment 	<ul style="list-style-type: none"> Primary: DLT, ORR Secondary: DOR, PFS, OS, Immune Response 	<ul style="list-style-type: none"> SSD: Jun. 2010 DE: 2019

Isatuximab (anti-CD38 mAb) Multiple Myeloma (MM)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
Lenalidomide Combination RRMM TCD11863 NCT01749969	Phase 1b Isatuximab, in Combination With lenalidomide and dexamethasone for the Tx of Relapsed or Refractory MM	60	<ul style="list-style-type: none"> Patients with diagnosis of MM and documentation of at least 2 prior therapies (induction therapy, autologous stem cell transplant, consolidation and maintenance therapy is considered one prior therapy) Open-label, Parallel assignment Isatuximab (escalating doses) + lenalidomide + dexamethasone Total duration for one patient: up to 21 days screening, at least 4 weeks Tx, up to 60 days follow-up 	<ul style="list-style-type: none"> Primary: N of patients with AE Secondary: ORR, PFS, PK, PD, Immunogenicity 	<ul style="list-style-type: none"> SSD: Feb. 2013 DE: 2019

Isatuximab (anti-CD38 mAb) Multiple Myeloma (MM)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
Pomalidomide Combination RRMM TCD14079 NCT02283775	Phase 1b Isatuximab, in combination with pomalidomide and dexamethasone for the Tx of Relapsed/Refractory MM	89	<ul style="list-style-type: none"> • Patients previously diagnosed with MM based on standard criteria and currently require Tx because MM has relapsed following a response • Open-label, Parallel assignment • Isatuximab + pomalidomide + dexamethasone • Part A, doses ranging for isatuximab, (5mg/kg, 10mg/kg, 20mg/kg); Part B isatuximab (10mg/kg) from a fixed infusion volume 	<ul style="list-style-type: none"> • Primary: DLTs, N of patients with AE • Secondary: ORR, PK, Immunogenicity, DOR, CB 	<ul style="list-style-type: none"> • SSD: May 2015 • DE: 2020

Isatuximab (anti-CD38 mAb) Multiple Myeloma (MM)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
Bortezomib Combination RRMM TCD13983 NCT02513186	Phase 1 Isatuximab in combination with bortezomib - based regimens in adult patients with newly diagnosed MM non eligible for transplantation	44	<ul style="list-style-type: none"> Patients with a diagnosis of MM with evidence of measurable disease, having received prior Tx with an IMiD and with at least 3 prior lines of therapy Open-label, Single Group assignment Isatuximab (escalating dose) + bortezomib + cyclophosphamide + dexamethasone: VCDI cohort (3-week screening, 50-week duration for induction and then up to disease progression, or unacceptable AEs + follow-up) Isatuximab + bortezomib + dexamethasone + lenalidomide: VRDI cohort to begin after VCDI completion (4-week screening, 24-week duration for induction and then up to disease progression, or unacceptable AEs, + follow-up) 	<ul style="list-style-type: none"> Primary: DLTs/VCDI For both VCDI & VRDI: ORR, CR Secondary: N of patients with AE, and significant changes in lab tests, PK, DOR 	<ul style="list-style-type: none"> SSD: Sep. 2015 DE: 1st set of data: 2018; next 2020, Full completion: 2023

Isatuximab (anti-CD38 mAb) Multiple Myeloma (MM)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
RRMM TED14154 NCT02514668	Phase 1 Safety, PK and Efficacy of isatuximab in patients with Relapsed/Refractory MM	64	<ul style="list-style-type: none"> Patients with a diagnosis of MM with evidence of measurable disease and with evidence of disease progression Open-label, Single Group assignment, isatuximab (escalating doses) Total duration for one patient: up to 21 days screening, Tx period up to disease progression or AEs , 60- day follow-up at least 	<ul style="list-style-type: none"> Primary: Part A: DLTs, N of patients with AE; Part B: ORR Secondary: PK, N of patients with AEs, DOR, CB, PFS, Immunogenicity 	<ul style="list-style-type: none"> SSD: Sep. 2015 DE: 2018

Isatuximab (anti-CD38 mAb) Multiple Myeloma (MM)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
ISLANDS (Japanese Patients) RRMM TED14095 NCT02812706	Phase 1 Phase 2 Isatuximab single-agent in Japanese patients with Relapsed and Refractory MM	42	<ul style="list-style-type: none"> Patients with a diagnosis of symptomatic MM, having received at least 3 prior lines of therapy OR whose disease is double refractory to an IMiD and a PI Open-label, Single Group assignment, isatuximab monotherapy Total duration for one patient: up to 21-day screening, Tx period up to disease progression or unacceptable AEs, post-Tx follow-up 	<ul style="list-style-type: none"> Primary: Phase 1: DLTs Phase 2: ORR Secondary: N of patients with AE, CB, OS, PFS, DOR, TTR, PK, PD, Immunogenicity 	<ul style="list-style-type: none"> SSD: Sep. 2016 DE: primary completion 2018; full completion 2019

Isatuximab (anti-CD38 mAb) Multiple Myeloma (MM)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
Cemiplimab Combination RRMM TCD14906 NCT03194867	Phase 1 Phase 2 Safety, PK and Efficacy of isatuximab in combination with cemiplimab in patients with Relapsed/Refractory MM	105	<ul style="list-style-type: none"> Patients with a diagnosis MM with evidence of measurable disease, having received prior Tx with an IMiD and with at least 3 prior lines of therapy Randomized, Open-label, Parallel Assignment Isatuximab + cemiplimab 3 Arms: Isa +cemi regimen 1; isa + cemi regimen 2; isa alone Total duration for one patient: up to 21-day screening, Tx period up to disease progression or unacceptable AEs, 3-month post-Tx follow-up. Cycle duration 28 days 	<ul style="list-style-type: none"> Primary: DLTs, N of patients with AE, ORR Secondary: CB, DOR, TTR, PFS, OS, PK, Immunogenicity (isatuximab and cemiplimab) 	<ul style="list-style-type: none"> SSD: Feb. 2018 DE: 2021

Isatuximab (anti-CD38 mAb) Multiple Myeloma (MM)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
ICARIA-MM RRMM EFC14335 NCT02990338	Phase 3 Isatuximab, pomalidomide, and dexamethasone to pomalidomide and dexamethasone in Refractory or Relapsed and RRMM	300	<ul style="list-style-type: none"> Isatuximab in combination with pomalidomide and low-dose dexamethasone, compared to pomalidomide and low-dose dexamethasone in patients with RRMM Randomized, Open-label, Parallel assignment 	<ul style="list-style-type: none"> Primary: PFS Secondary: ORR, OS, TTP, PFS, DOR 	<ul style="list-style-type: none"> SSD: Jan. 2017 DE (1st Part)⁽¹⁾: 2018; full completion 2020

Isatuximab (anti-CD38 mAb) Multiple Myeloma (MM)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
IKEMA RRMM EFC15246 NCT03275285	Phase 3 Isatuximab combined with carfilzomib and dexamethasone vs. carfilzomib with dexamethasone in patients With Relapse and/or Refractory MM previously treated with 1 to 3 prior lines	300	<ul style="list-style-type: none"> Patients with MM previously treated with prior 1 to 3 lines and with measurable serum M-protein (≥ 0.5 g/dL) and/or urine M-protein (≥ 200 mg/24 hours) Randomized, Open-label, Parallel assignment, 2-arm: (a) isatuximab +carfilzomib+dexamethasone, (b) carfilzomib+dexamethasone 	<ul style="list-style-type: none"> Primary: PFS Secondary: ORR, % of patients with CR, and VGPR, OS, TTP, Second PFS, DOR, AE, PK, Immunogenicity 	<ul style="list-style-type: none"> SSD: Oct. 2017 DE (1st Part)⁽¹⁾: 2020; full completion: 2023

Isatuximab (anti-CD38 mAb) Multiple Myeloma (MM)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
IMROZ NDMM EFC12522 NCT03319667	Phase 3 Isatuximab in combination with bortezomib (Velcade®), lenalidomide (Revlimid®) and dexamethasone vs. bortezomib, lenalidomide and dexamethasone in patients with newly diagnosed MM not eligible for transplant	440	<ul style="list-style-type: none"> Newly diagnosed MM not eligible for transplant due to age (≥ 65 years) or patients < 65 years with comorbidities impacting possibility of transplant or patient's refusal of transplant Randomized, Open-label, Parallel assignment IVRd arm (Isatuximab/bortezomib/lenalidomide/dexamethasone) VRd arm (Bortezomib/lenalidomide/dexamethasone) Ird crossover arm (Isatuximab/lenalidomide/dexamethasone) Total duration for each patient: screening period up to 4 weeks, induction period of 24 weeks, continuous Tx period and crossover when applicable 	<ul style="list-style-type: none"> Primary: PFS Secondary: ORR, % of patients with CR, and VGPR, OS, TTP, DOR, PFS on next line of therapy (PFS2), AE, PK, Immunogenicity, QOL 	<ul style="list-style-type: none"> SSD: 2017 DE (1st Part)⁽¹⁾: 2022

Isatuximab (anti-CD38 mAb) combination cemiplimab (PD-1 inhibitor) – Advanced Malignancies

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
Advanced Malignancies ACT15319 NCT03367819	Phase 1/2 Safety and tolerability of Isatuximab in combination with cemiplimab in patients with metastatic castration-resistant prostate cancer (mCRPC) or patients with non-small cell lung cancer (NSCLC)	134	<ul style="list-style-type: none"> In Patients with metastatic, castration-resistant prostate cancer (mCRPC) who are naïve to anti-programmed cell death-1 (PD-1)/programmed cell death-ligand 1 (PDL-1)-containing therapy, or non-small cell lung cancer (NSCLC) who progressed on anti-PD-1/PDL-1-containing therapy, Randomized, Open-Label, Parallel Assignment Isatuximab alone or in combination with cemiplimab Total duration per patient up to 28 months including 28 days screening period, , up to 24 months ttmt period and 3 months safety FU 	<ul style="list-style-type: none"> Primary: Safety, tolerability, RR Secondary: Immunogenicity (isa and cemi), PK, tumor burden change, DR, PFS, 	<ul style="list-style-type: none"> SSD: 2018 DE: 2021

Cemiplimab (PD-1 inhibitor) Advanced Malignancies (AM)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
AM R2810-ONC-1423 NCT02383212	Phase 1 A first-in-human study of repeat dosing with cemiplimab, as single therapy and in combination with other Anti-Cancer therapies in patients with AM	398	<ul style="list-style-type: none"> • Non-Randomized, Open-label, Parallel assignment, ascending-dose • Monotherapy, cemiplimab alone • Dual combination: cemiplimab in combination with hypofractionated radiotherapy or with cyclophosphamide or with docetaxel • Triple combination: cemiplimab with hypofractionated radiotherapy plus cyclophosphamide, or hypofractionated radiotherapy plus GM-CSF or carboplatin plus paclitaxel or carboplatin plus pemetrexed or carboplatin plus docetaxel • Quadruple combination: cemiplimab with hypofractionated radiotherapy plus GM-CSF plus cyclophosphamide 	<ul style="list-style-type: none"> • Primary: TEAE, Incidence of abnormal laboratory findings, N of participants with DLT • Secondary, RECIST as measured by CT or MRI, Immune-Related Response 	<ul style="list-style-type: none"> • SSD: Jan. 2015 • DE: 2020

Cemiplimab (PD-1 inhibitor) Advanced Malignancies (AM)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
PK in Japanese patients AM R2810-ONC-1622 NCT03233139	Phase 1 To investigate the safety and PKs of cemiplimab monotherapy in Japanese patients with AM	14	<ul style="list-style-type: none"> Histologically or cytologically confirmed diagnosis of malignancy with no alternative standard-of-care therapeutic option Single Group assignment, Open-label 	<ul style="list-style-type: none"> Primary: TEAEs cemiplimab PK parameters Secondary: Immunogenicity against cemiplimab 	<ul style="list-style-type: none"> SSD: Sep. 2017 DE: primary 2019; full completion 2023

Cemiplimab (PD-1 inhibitor) Combination REGN3767 Advanced Malignancies (AM)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
R3767-ONC-1613 NCT03005782	Phase 1 To investigate the safety and PKs of REGN3767 (anti LAG-3 mAb) to determine the recommended Phase 2 dose (RP2D) as monotherapy and in combination with cemiplimab in patients with advanced malignancies	546	<ul style="list-style-type: none"> Histologically or cytologically confirmed diagnosis of malignancy with no alternative standard-of-care therapeutic option Non-randomized, Parallel Group assignment, Open-label Group A: REGN3767, 4 sequential dose cohorts, each cohort receiving 1 of 3 ascending dose levels. 1 tumor-specific cohort treated with the RP2D during dose expansion Group B: REGN3767+cemiplimab, same design; 9 tumor-specific cohorts treated with RP2D 	<ul style="list-style-type: none"> Primary: DLTs, PK parameters, AEs, SAEs, death and lab. abnormalities, response rate Secondary: Response rate, duration of response, disease control rate, PFS, Aes, SAEs, death, lab. abnormalities immunogenicity 	<ul style="list-style-type: none"> SSD: Nov. 2016 DE: Primary completion 2021, full completion 2022

Cemiplimab (PD-1 inhibitor) Melanoma

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
Biomarkers Melanoma R2810-ONC-1606 NCT03002376	Phase 1 Exploratory Tumor Biopsy-driven study to understand the relationship between biomarkers and clinical response in Melanoma patients receiving cemiplimab	30	<ul style="list-style-type: none"> For Histologically confirmed diagnosis of stage III (unresectable) or stage IV melanoma with at least 1 lesion that is measurable by RECIST 1.1 criteria and accessible for biopsies Non-Randomized, Open-label, Parallel assignment Group 1: Patients with metastatic CSCC: to distant sites or lymph nodes. cemiplimab administered intravenously every 2 weeks Group 2: Patients with unresectable locally advanced CSCC. cemiplimab administered intravenously every 2 weeks Group 3: Patients with metastatic CSCC, to distant sites or lymph nodes. cemiplimab administered intravenously every 3 weeks 	<ul style="list-style-type: none"> Primary: Correlation between changes in the tumor microenvironment and the change in tumor volume following cemiplimab Tx Secondary: Correlation between baseline tumor characteristics and the change in tumor volume following Tx, cemiplimab serum concentrations, antibodies levels, PFS, ORR 	<ul style="list-style-type: none"> SSD: Apr. 2017 DE (1st Part)⁽¹⁾: 2018

Cemiplimab (PD-1 inhibitor) Head and Neck

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
Biomarkers Head & Neck R2810-ONC-1655 NCT03198130	Phase 1 Exploratory Tumor Biopsy-driven study to understand the relationship between biomarkers and clinical response in Immunomodulatory Treatment-Naïve patients with Recurrent and/or Metastatic Squamous Cell Carcinoma of Head and Neck receiving cemiplimab	30	<ul style="list-style-type: none"> For Histologically confirmed diagnosis recurrent and/or metastatic SCCHN (squamous cell carcinoma of the head and neck) with no curative options with at least 1 lesion that is measurable by Response Evaluation Criteria in Solid Tumors (RECIST) Open-label, Single Group Assignment 	<ul style="list-style-type: none"> Primary: Correlation between changes in the tumor microenvironment and the change in tumor volume following cemiplimab Tx Secondary: Correlation between baseline tumor characteristics and the change in tumor volume following Tx, ORR, PFS, TAES, cemiplimab serum concentration, anti-cemiplimab antibodies level 	<ul style="list-style-type: none"> SSD: Jul. 2017 DE (1st Part) ⁽¹⁾: 2019; full completion 2020

Cemiplimab (PD-1 inhibitor) Cutaneous Squamous Cell Carcinoma (CSCC)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
Advanced CSCC R2810-ONC-1540 NCT02760498	Phase 2 Cemiplimab monotherapy for patients with metastatic (nodal or distant) CSCC (Groups 1 and 3) or with unresectable locally advanced CSCC (Group 2)	182	<ul style="list-style-type: none"> Non-Randomized, Open-label, Parallel assignment Group 1: Patients with metastatic CSCC: to distant sites or lymph nodes cemiplimab administered intravenously every 2 weeks Group 2: Patients with unresectable locally advanced CSCC. cemiplimab administered intravenously every 2 weeks Group 3: Patients with metastatic CSCC: to distant sites or lymph nodes, cemiplimab administered intravenously every 3 weeks 	<ul style="list-style-type: none"> Primary: ORR (96 weeks), Groups 1 and 3: RECIST version 1.1 will be used to determine ORR, Group 2: Clinical response criteria will be used to determine ORR Secondary: Investigator Assessments of ORR, DOR, PFS, OS, CRR 	<ul style="list-style-type: none"> SSD: May 2016 DE: 2020
Expanded Access CSCC R2810-ONC-17103 NCT03492489	Expanded Access Tx IND/Protocol Provide access to cemiplimab to patients with mCSCC or locally advanced CSCC, who are not candidate for surgery prior to cemiplimab being commercially available	Intermediate-size Population			

Cemiplimab (PD-1 inhibitor) Basal Cell Carcinoma (BCC)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
BCC R2810-ONC-1620 NCT03132636	Phase 2 Cemiplimab in patients with Advanced BCC who experienced progression of disease on Hedgehog Pathway Inhibitor Therapy, or were intolerant of Prior Hedgehog Pathway Inhibitor Therapy	137	<ul style="list-style-type: none"> Patients with confirmed diagnosis of invasive BCC Non-Randomized, Open-label, Parallel assignment Group 1: Patients with metastatic BCC Group 2: Patients with unresectable locally advanced BCC 	<ul style="list-style-type: none"> Primary: ORR for mBCC measured by RECIST version 1.1 ORR for unresectable locally advanced BCC measured by Composite Response Criteria Secondary: DOR, CR, PFS, OS 	<ul style="list-style-type: none"> SSD: July 2017 DE (1st Part) ⁽¹⁾: 2018; full completion 2020

Cemiplimab (PD-1 inhibitor) Non-Small Cell Lung Cancer (NSCLC)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
mNSCLC R2810-ONC-1624 NCT03088540	Phase 3 First-line Tx in patients with advanced or metastatic NSCLC whose tumors express PD-L1, vs. Platinum Based Chemotherapy	300	<ul style="list-style-type: none"> For histologically or cytologically documented squamous or non squamous NSCLC with stage IIIB or stage IV disease who received no prior systemic Tx for recurrent or metastatic NSCLC Randomized, Open-label, Cross-over assignment Active Comparator: Standard-of-care chemotherapy: paclitaxel + cisplatin OR paclitaxel + carboplatin OR gemcitabine + cisplatin or gemcitabine + carboplatin OR Pemetrexed + cisplatin followed by optional pemetrexed maintenance OR pemetrexed + carboplatin followed by optional pemetrexed maintenance 	<ul style="list-style-type: none"> Primary: PFS as assessed by a blinded Independent review committee using RECIST 1.1 Secondary: OS, Objective response rates, BOR, DOR 	<ul style="list-style-type: none"> SSD: May 2017 DE: 2021

Cemiplimab (PD-1 inhibitor) Non-Small Cell Lung Cancer (NSCLC)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
mNSCLC R2810-ONC-16113 NCT03409614	Phase 3 Combination of cemiplimab, ipilimumab and Platinum-based Doublet Chemotherapy in 1 st Line Tx of patients with advanced or metastatic NSCLC with tumors expressing PD-L1<50%	690	<ul style="list-style-type: none"> For histologically or cytologically documented squamous or non squamous NSCLC with stage IIIB disease who are not candidates for Tx with definitive concurrent chemoradiation or stage IV disease who received no prior systemic Tx for recurrent or metastatic NSCLC Randomized, Open-label, Parallel assignment Arm 1: Standard of care (SOC) Arm 2: cemiplimab + SOC Arm 3: cemiplimab + abbreviated chemotherapy + ipilimumab 	<ul style="list-style-type: none"> Primary: PFS as assessed by a blinded Independent review committee using RECIST 1.1 Secondary: OS, ORR, TEAEs, DLTs, SAEs, death, lab. abnormalities, OS, QoL 	<ul style="list-style-type: none"> SSD: Mar. 2018 DE: 2022

Cemiplimab (PD-1 inhibitor) Non-Small Cell Lung Cancer (NSCLC)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
mNSCLC R2810-ONC-1763 NCT03430063	Phase 2 Combination of standard and High dose of cemiplimab and ipilimumab in 2 nd Line Tx of patients with mNSCLC with tumors expressing PD-L1 < 50%	201	<ul style="list-style-type: none"> For histologically or cytologically documented squamous or non squamous NSCLC with stage IIIB and not candidates for definitive chemoradiation or stage IV. Patients immunotherapy naïve and having received one prior cytotoxic regimen. Randomized, Open-label, Parallel assignment Arm 1: cemiplimab standard dose Arm 2: cemiplimab + ipilimumab standard doses Arm 3: cemiplimab High dose 	<ul style="list-style-type: none"> Primary: ORR Secondary: OS, PFS, TEAEs, SAEs, death, lab. abnormalities, QoL, immunogenicity, hair pigmentation, tumor burden, tumor volume, PK, markers, 	<ul style="list-style-type: none"> SSD: May 2017 DE: 2021

Cemiplimab (PD-1 inhibitor) Combination REGN4659 Non-Small Cell Lung Cancer (NSCLC)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
mNSCLC R4659-ONC-1795 NCT03580694	Phase 1 Cemiplimab in combination with REGN4959 in the Tx of patients with advanced or mNSCLC	134	<ul style="list-style-type: none"> For histologically or cytologically documented squamous or non squamous NSCLC with stage IIIB or stage IV who received no prior systemic Tx for recurrent or mNSCLC; with expression of PDL-1in $\geq 50\%$ of tumors cells Non-Randomized, Open-label, Parallel assignment Arm 1: cemiplimab Arm 2: cemiplimab + REGN4659 	<ul style="list-style-type: none"> Primary: DLTs, TEAEs, immune-related AEs, SAEs, deaths, lab. abnormalities, ORR, PK of both products Secondary: ORR, BOR, DOR, disease control rate, PFS, OS, 	<ul style="list-style-type: none"> SSD: June 2018 DE: 2021

Cemiplimab (PD-1 inhibitor) Cervical cancer (CC)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
CC R2810-ONC-1676 NCT03257267	Phase 3 Cemiplimab vs. therapy of IC chemotherapy in Recurrent or Metastatic Platinum-Refractory CC	436	<ul style="list-style-type: none"> Patients with recurrent or metastatic platinum-refractory CC treated with either REGN2810 or IC chemotherapy Randomized, Open-label, Parallel assignment, Tx cycle 6 weeks, Planned Tx for up to 96 weeks 	<ul style="list-style-type: none"> Primary: OS Secondary: PFS, ORR, DOR, QOL 	<ul style="list-style-type: none"> SSD: Oct. 2017 DE (1st Part) ⁽¹⁾: 2020; next 2022; full completion 2023

Cemiplimab (PD-1 inhibitor) Combination REGN4018 Ovarian cancer (OC)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
CC R4810-ONC-1721 NCT03564340	Phase 1/2 REGN4018 alone or in combination with cemiplimab in patients with Platinum-Resistant Ovarian Cancer	264	<ul style="list-style-type: none"> Histologically or cytologically confirmed diagnosis of advanced, epithelial ovarian (except carcinosarcoma), primary peritoneal, or fallopian tube cancer with CA-125 \geq 2 xULN, progression or relapse within 6 months of the most recent Tx with Platinum-containing chemotherapy, documented progression and no standard therapy options Non- Randomized, Open-label, Parallel assignment, Arm 1: REGN4018 Arm 2: REGN4018 + cemiplimab 	<ul style="list-style-type: none"> Primary: DLTs, TEAEs, SAEs, deaths, lab abnormalities, drugs serum concentrations, ORR Secondary: BOR, DOR, disease control, PFS, CA-125 	<ul style="list-style-type: none"> SSD: May 2018 DE: 2022

SAR439459 (TGFβ inhibitor mAb)

Advanced Solid Tumors (AST)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
AST Monotherapy and combination with cemiplimab TCD14678 NCT03192345	Phase 1/1b PK, PD and Anti-tumor activity of SAR439459 Monotherapy and in combination with cemiplimab in adult patients with AST	130 expected	<ul style="list-style-type: none"> • Patients with histologically confirmed, advanced unresectable or metastatic solid tumor • Randomized, Open-label, Parallel assignment • Part 1A: SAR439459 monotherapy escalating doses/14-day cycle • Part 2A: SAR439459 monotherapy/14-day cycle with the previously recommended dose • Part 1B: SAR439459 escalating dose + cemiplimab standard dose /14-day cycle • Part 2B: SAR439459 at previously recommended dose + cemiplimab standard dose / 14-day • Escalation periods non randomized followed par expansion periods randomized 	<ul style="list-style-type: none"> • Primary: DLTs (Part 1), ORR (Part 2) • Secondary: Safety, Immunogenicity, PFS, TTP, PK 	<ul style="list-style-type: none"> • SSD: Jun. 2017 • DE: 2021

SAR408701 (maytansin loaded anti-CEACAM5 mAb) Advanced Solid Tumors (AST) 1/2

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
First-in-Human TED13751 NCT02187848	Phase 1 Phase 2 PK and antitumor activity of SAR408701 in patients with AST	233 expected	<ul style="list-style-type: none"> Patients with locally advanced or metastatic solid malignant tumor Non-Randomized, Open-label, Parallel assignment Arm 1 : SAR408701 monotherapy escalating cohorts Arm 2: SAR408701 expansion cohort in CRC with MTD previously defined Arm 3: SAR408701 expansion cohort lung adenocarcinoma at MTD Arm 4: SAR408701 expansion cohort gastric adenocarcinoma at MTD Arm 5: SAR408701 loading dose at first cycle followed by MTD Arm 6: SAR408701 expansion cohort lung adenocarcinoma (Lung bis) at MTD Arm 7: SAR408701 expansion cohort colorectal cancer (CRC-L) at MTD 	<ul style="list-style-type: none"> Primary: MTD, Anti-tumor response RECIST Secondary: Safety, Immunogenicity, PK 	<ul style="list-style-type: none"> SSD: Sep. 2014 DE: 2019

SAR408701 (maytansin loaded anti-CEACAM5 mAb) Advanced Solid Tumors (AST) 2/2

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
Japanese patients Monotherapy and Combination TCD15054 NCT03324113	Phase 1 Safety and PK of SAR408701 Monotherapy and in combination with other anti-tumor drug in Japanese patients with Advanced Malignant Solid Tumors	27	<ul style="list-style-type: none"> • Patients with malignant solid tumor • Non-Randomized, Open-label, Sequential assignment • Phase 1 : SAR408701 monotherapy escalating doses/ 4 weeks • Phase 1B: SAR408701 at MTD in combinations with other anti-tumor drugs, 4 weeks 	<ul style="list-style-type: none"> • Primary: DLTs, Phase 1 and 1B • Secondary: Safety, Immunogenicity, PK, Plasma CEACAM5 levels, Anti-tumor response RECIST 	<ul style="list-style-type: none"> • SSD: Oct. 2017 • DE: 2020

SAR439859 (SERD)

Breast cancer

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
TED14856 NCT03284957	Phase 1 Phase 2 SAR439859 single agent and in combination with palbociclib in Postmenopausal Women with Estrogen Receptor Positive Advanced Breast Cancer	156	<ul style="list-style-type: none"> • Non-Randomized, Open-label, Parallel Assignment • Part A: SAR439859 monotherapy dose escalation, • Part C: dose escalation for the combination SAR439859 and palbociclib, • Part B: SAR439859 dose expansion from the dose determined in part A, • Part D: combination SAR439859 and palbociclib at the doses recommended from part B • Sar439859 administered in 28-day cycle; palbociclib in 21-day cycle 	<ul style="list-style-type: none"> • Primary: Parts A & C:DLTs, Parts B & D: ORR • Secondary: Safety, ORR, DCR, DR, PK for both drugs, CYP450 3A induction/inhibition, ER occupancy/PET imaging 	<ul style="list-style-type: none"> • SSD: Sept. 2017 • DE: 2020

GZ402666 (avalglucosidase alfa)

Pompe disease (PD) 1/3

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
COMET Late Onset EFC14028 NCT02782741	Phase 3 To compare efficacy and safety of Enzyme Replacement therapies avalglucosidase alfa (neoGAA) and alglucosidase alfa (Myozyme®/Lumizyme®) in Tx-naïve patients with Late-onset PD	96	<ul style="list-style-type: none"> Repeated Biweekly Infusions of avalglucosidase alfa (GZ402666) and alglucosidase alfa in Tx-naïve patients with late-onset PD age 3 years and older Randomized, Double-Blind, Parallel Assignment Total study duration for one patient: 3 years [14-day screening, 49-week blinded Tx period, 96-week open-label Tx and 4-week post-Tx observation period 	<ul style="list-style-type: none"> Primary: Change in percent predicted forced vital capacity (%FVC) in the upright position, from baseline to 12 months Secondary: Change from baseline to 12 months in six-minute walk test distance walked, maximal inspiratory / expiratory pressure (% predicted), hand-held dynamometry measurement of lower extremity muscle strength in Quick Motor Function Test scores, and 12- Item Short-form health survey scores 	<ul style="list-style-type: none"> SSD: Oct. 2016 DE (1st Part): 2020

GZ402666 (avalglucosidase alfa)

Pompe disease (PD) 2/3

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
Mini-COMET Infantile Onset ACT14132 NCT03019406	Phase 2 To assess safety and efficacy of avalglucosidase alfa (neoGAA) in Pediatric patients with infantile-onset PD previously treated With alglucosidase alfa (Myozyme®/Lumizyme®)	20	<ul style="list-style-type: none"> Repeated bi-weekly infusions of avalglucosidase alfa In Patients with Infantile-onset PD previously treated with alglucosidase alfa (Myozyme®/Lumizyme®) who demonstrate clinical decline or sub-optimal clinical response Randomized, Open-label, Ascending dose, Parallel assignment Total study duration for one patient: 3 years [14-day screening, 25-week Tx period, a 120-week extension period and 4-week post-Tx observation period 	<ul style="list-style-type: none"> Primary: N of participants with AE Secondary: PK parameters, Change at 6 months from baseline in Gross Motor Function (GMF) Measure-88 Test, revised GMF Classification System score, Pompe specific Pediatric Evaluation of Disability Inventory, Functional Skills Scale, Mobility Domain Test score and Quick Motor Function Test scores, Left Ventricular Mass Index, Eyelid position measurements, Creatine kinase value 	<ul style="list-style-type: none"> SSD: Oct. 2017 DE (1st Part): 2022

GZ402666 (avalglucosidase alfa)

Pompe disease (PD) 3/3

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
NEO-EXT LTS13769 NCT02032524	Phase 2 Phase 3 Long-term safety and PK of repeated biweekly infusions of avalglucosidase alfa (neoGAA) in patients with PD	21	<ul style="list-style-type: none"> Repeated biweekly infusions of avalglucosidase alfa in patients with PD who previously completed a avalglucosidase alfa study [adult, senior] Non-randomized, Open-label, Parallel assignment Total study duration for one patient: 6 years [until the patient withdraws, the Investigator withdraws the patient, or the Sponsor terminates the study] 	<ul style="list-style-type: none"> Primary: AEs and TEAEs, including IARs & deaths, Hematology, biochemistry and urinalysis, vital signs Secondary: ECG, PK parameters, anti-avalglucosidase alfa antibodies, and neutralizing antibody formation in anti-avalglucosidase alfa positive patients, anti-avalglucosidase alfa IgG antibodies, Skeletal muscle glycogen content, Qualitative and quantitative muscle degenerative assessments MRI, Urinary Hex4, plasma analyses of circulating mRNA and micro RNA, Serum analyses of skeletal muscle RNA expression 	<ul style="list-style-type: none"> SSD: Feb. 2014 DE: 2020

Olipudase Alfa (rhASM ERT) 1/3 Acid Sphingomyelinase Deficiency (ASMD)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
ASCEND Niemann-Pick disease type B⁽¹⁾ DFI12712 NCT02004691	Phase 2 Phase 3 Efficacy, Safety, PD, and PK study of olipudase alfa in patients with ASD	36	<ul style="list-style-type: none"> Randomized, Double-blinded, Placebo-controlled, Parallel assignment Total study duration for one patient at least 3 years up to 5 years and 3 months [2-month screening, 52-week double-blind Tx period, 4-year and 1 month open label extension period with olipudase 	<ul style="list-style-type: none"> Primary: % change in spleen volume, % change in diffusing capacity of the lung for carbon monoxide (DLco) Secondary: Change in splenomegaly-related symptom score (except US, where it is part of the primary "combination spleen endpoint"), % change in liver volume, % change in platelet count, Change in fatigue severity as measured by item 3 of the Brief Fatigue Inventory scale, Change in pain severity as measured by item 3 of the Brief Pain Inventory scale, Change in dyspnea severity as measured by the Functional Assessment of Chronic Illness Therapy dyspnea tool 	<ul style="list-style-type: none"> SSD: June 2016 DE (1st Part)⁽²⁾: 2019

Olipudase Alfa (rhASM ERT) 2/3 Acid Sphingomyelinase Deficiency (ASMD)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
ASCEND Peds DFI13803 NCT02292654	Phase 1 Phase 2 Safety, Tolerability, PK, and efficacy evaluation of ollipudase alfa in pediatric patients <18 years of age with ASMD	20	<ul style="list-style-type: none"> Open-label, ascending dose, Single group assignment Total study duration for one patient approximately 18 months [up to 60-day screening, 64-week Tx period, 37-day post Tx period except if patient enrolled in a long-term extension study] 	<ul style="list-style-type: none"> Primary: safety parameters and Clinically significant changes in laboratory parameters, Clinically significant changes in physical examinations Secondary: PK parameters, Change in sphingomyelin levels and sphingomyelin metabolite levels 	<ul style="list-style-type: none"> SSD: June 2015 DE: 2019

Olipudase Alfa (rhASM ERT) 3/3

Acid Sphingomyelinase Deficiency (ASMD)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
Long-Term LTS13632 NCT02004704	Phase 2 Long-term study of olipudase alfa in patients with ASMD	17	<ul style="list-style-type: none"> For patients who have completed a previous study with olipudase alfa (DFI13803 for pediatric patients, and DFI13412 for adult patients) Open-label, Single group assignment Total study duration for one patient: 5 years 	<ul style="list-style-type: none"> Primary: Safety parameters and physical examinations including neurologic examinations, clinical laboratory tests, inflammatory biomarkers, immune response assessment, vital signs, echocardiogram and electrocardiogram, liver biopsy and liver ultrasound/doppler for patients previously enrolled in DFI13412. Secondary: Spleen and Liver Volumes, Pulmonary imaging and function tests, Hematology and Lipid profiles, Health Outcomes Questionnaires For pediatrics patients: Hand X-ray for bone age and bone maturation, Tanner Staging and Linear patient growth by height Z-score 	<ul style="list-style-type: none"> SSD: Dec. 2013 DE: 2023

Venglustat (GCS inhibitor) Fabry disease (FD)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
FABRY LONG-TERM LTS14116 NCT02489344	Phase 2 Long-term safety, PD, and exploratory efficacy of venglustat in Tx-naïve adult male patients with FD	8	<ul style="list-style-type: none"> Male patients with FD who previously completed study ACT13739 Open-label, Single group Assignment Total study duration for one patient: up to 31 months 	<ul style="list-style-type: none"> Primary: Safety profile, Clinically significant changes in laboratory parameter, and physical examinations Secondary: Change from baseline in plasma globotriaosylceramide (GL-3), plasma lyso GL-3, Change from baseline in plasma glucosylceramide (GL 1), Urine GL-3 	<ul style="list-style-type: none"> SSD: Jul. 2015 DE: 2018

Venglustat (GCS inhibitor) Gaucher disease (GD) Type 3

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
LEAP GD Type 3 PDY13949 NCT02843035	Phase 2 Tolerability, PK, PD, and exploratory efficacy of venglustat in combination with cerezyme in adult patients with GD Type 3	10	<ul style="list-style-type: none"> 156-week Three part, Open-label, Single group Assignment Part 1: Evaluate CNS biomarkers in adult GD type 3 patients that distinguish GD3 from GD type 1, Screen adult GD3 patients who qualify for Ttmt with venglustat in Part 2, Total duration 45 days Part 2 and 3: Safety and tolerability in GD3 patients, Total duration up to 156 weeks including 2 part-ttmt of 52 weeks (Part 2) and 104 weeks (Part 3) for long term follow-up, respectively 	<ul style="list-style-type: none"> Primary: N of patients with AE, assessment of PD parameters (GL-1 and lyso GL1) in CSF and plasma Secondary: PK parameters (CSF and Plasma) 	<ul style="list-style-type: none"> SSD: Jan. 2017 DE ^(1st Part)(1): 2021

Venglustat (GCS inhibitor)

Autosomal Dominant Polycystic Kidney Disease (ADPKD)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
ADPKD EFC15392 NCT03523728	Phase 3 Efficacy, Safety, Tolerability and PK of venglustat in patients at risk of rapidly progressive ADPKD	560	<ul style="list-style-type: none"> Randomized, double-blind, placebo-controlled 2-stage study (18 and 24 months) Study duration per participant is 26 months (maximal) per stage, including a screening period of 15 days, run-in period of 2 weeks, a 24-month treatment period, and a follow-up 1 month after final dose 	<ul style="list-style-type: none"> Primary: Rate of change in total kidney volume (TKV) based on magnetic resonance imaging (MRI) and rate of change in glomerular filtration rate (eGFR) Secondary: Rate of change in eGFR, rate of change in TKV, rate of change in urine osmolaity, rate of change in nocturia, adverse events, assessment of PK, change in lens clarity 	<ul style="list-style-type: none"> SSD: Aug. 2018 (FPI in Q4 2018) DE (1st Part)⁽¹⁾: 2021

Eliglustat

Gaucher's Disease (GD) 1/2

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
ELIKIDS GD Type 1/ Type 3 EFC13738 NCT03485677	Phase 3 PK, efficacy and safety with or without Imiglucerase in pediatric patients with GD Type I/Type 3	60	<ul style="list-style-type: none"> Non-randomized, open label, two cohort (with and without imiglucerase) Cohort 1: eliglustat monotherapy Cohort 2: eligustat plus imiglucerase 	<ul style="list-style-type: none"> Primary: PK (Cmax and AUC), adverse events Secondary: changes from baseline as absolute change in g/dL for hemoglobin, % change for platelets, liver volume, and spleen volume; improvement in pulmonary disease, improvement in bone disease, thrombocytopenia, and quality of life 	<ul style="list-style-type: none"> SSD: Apr. 2018 DE (1st Part)⁽¹⁾: 2022

Eliglustat

Gaucher's Disease (GD) 2/2

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
EXOSKEL GD Type 1 EFC13781 NCT02536755	Phase 3 Long Term skeletal response to eliglustat in GD Type 1 adult patients who successfully completed Phase 2 or phase 3 studies	32	<ul style="list-style-type: none"> Single group assignment, open label 	<ul style="list-style-type: none"> Primary: change from baseline in bone marrow infiltration, bone mineral density (hips and lumbar spine), skeletal imaging GD bone disease manifestations (lytic lesions, osteonecrosis, fractures and infarcts), clinical GD manifestations (mobility, bone pain, bone crisis), and bone biomarkers Secondary: quality of life, measurement of GD Type 1 biomarkers and safety (i.e. incidence of adverse events, change from baseline in laboratory assessments (hematology), physical examinations) 	<ul style="list-style-type: none"> SSD: Oct. 2015 DE (1st Part)⁽¹⁾: 2021

Teriflunomide

Multiple Sclerosis (MS)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
TERIKIDS RMS EFC11759 NCT02201108	Phase 3 Efficacy, Safety and PK of teriflunomide in Pediatric Patients With Relapsing Forms of MS	165	<ul style="list-style-type: none"> Patients with RMS meeting the criteria of MS based on McDonald criteria 2010 and International Pediatric MS Study Group criteria for pediatric MS With at least one relapse (or attack) in the 12 months preceding randomization or at least two relapses (or attack) in the 24 months preceding randomization Randomized, Double-Blind, Placebo-Controlled, Parallel Group , Tx 96 weeks followed by Open-label extension (96 weeks up to a max of 192 weeks after randomization), follow-up 4 weeks after Tx discontinuation 	<ul style="list-style-type: none"> Primary: Time to first clinical relapse after randomization Secondary: % of relapse free patients, N of new/newly enlarged T2 lesions, N of T1 Gd-enhancing T1 lesions , Change in volume of T2 lesions , of T1 hypointense lesions , brain atrophy, % of patients free of new or enlarged MRI T2-lesions, Change in performance on SDMT and Cognitive Battery Test , Safety, PK 	<ul style="list-style-type: none"> SSD: Jul. 2014 DE: 2019

Alemtuzumab

Relapsing Remitting Multiple Sclerosis (RRMS)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
LemKids RRMS EFC13429 NCT03368664	Phase 3 Efficacy, Safety and Tolerability of Alemtuzumab in Pediatric Patients With Relapsing Remitting MS (RRMS) with disease activity on prior disease modifying therapy DMT	50	<ul style="list-style-type: none"> In pediatric patients from 10 to <18 years of age with RRMS with disease activity on prior DMT. Open-label, rater-blinded, single-arm, cross-over study The study will consist of different phases: <ul style="list-style-type: none"> Prior DMT Phase (~4 months) – efficacy measurements on current DMT Alemtuzumab Treatment Phase (~2 years) - The MRI based primary efficacy endpoint will be assessed over a 4 month period during this phase compared to an equal period during the prior DMT phase Safety Monitoring Phase – safety monitoring for all patients treated with alemtuzumab (4 years post last treatment with alemtuzumab) 	<ul style="list-style-type: none"> Primary: The number of new or enlarging T2 lesions on brain MRI, during continuation of prior DMT (Period 1) compared to an equal period after the first course of alemtuzumab treatment (Period 2) Secondary: The proportion of patients with new or enlarging T2 lesions , Annualized relapse rate at Year 2, Assessment of cognition test scores, Additional secondary endpoints, including PK/PD parameters and Quality of Life (QoL) measures. 	<ul style="list-style-type: none"> SSD: Oct. 2017 DE: 2025

SAR422459 (ABCA4 gene therapy) Stargardt Disease

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
Stargardt's Macular Degeneration TDU13583 NCT01367444	Phase 1 Phase 2/2a Safety and tolerability of ascending doses of SAR422459 in patients with Stargardt's Macular Degeneration	46	<ul style="list-style-type: none"> Patients with a diagnosis of Stargardt's Macular Degeneration, with at least one pathogenic mutant ABCA4 allele on each chromosome Non-randomized, Single Group assignment, Open-label, ascending doses 	<ul style="list-style-type: none"> Primary: IAE, Change from baseline in ocular safety assessments Secondary: Delay in retinal degeneration 	<ul style="list-style-type: none"> SSD: Jun. 2011 DE: 2020
Stargardt's Macular Degeneration LTS13588 SG1/002/11 NCT01736592	Phase 1/2 Follow-up study of SAR422459 in patients With Stargardt 's Macular Degeneration	46	<ul style="list-style-type: none"> Long Term safety and tolerability of SAR422459 in patients with Stargardt 's Macular Degeneration No ttmt administered, in this LTS only follow-up after ttmt in TDU13583 Patients will be followed for 15 years after treatment 	<ul style="list-style-type: none"> Primary: IAE Secondary: Delay in retinal degeneration 	<ul style="list-style-type: none"> SSD: Dec. 2012 DE: 2034

SAR421869 (Myosin 7A gene therapy) Usher 1B Syndrome

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
UshStat® Usher Syndrome Type 1B TDU13600 NCT01505062	Phase 1 Phase 2a Safety and tolerability of ascending doses of subretinal injections of UshStat® in patients with Retinitis Pigmentosa associated with Usher syndrome Type 1B	18	<ul style="list-style-type: none"> Patients with clinical and molecular diagnosis of Retinitis Pigmentosa associated with Usher Syndrome type 1B. With at least one pathogenic mutation in the MYO7A gene on each chromosome Non-randomized, Single Group assignment, Open-label, ascending doses 	<ul style="list-style-type: none"> Primary: IAE Secondary: Delay in retinal degeneration 	<ul style="list-style-type: none"> SSD: Mar. 2012 DE: 2021
UshStat® Usher Syndrome Type 1B LTS13619 NCT02065011	Phase 2b Long-Term Safety, Tolerability and Biological Activity of UshStat® in Patients With Usher Syndrome Type 1B	18	<ul style="list-style-type: none"> Long-term follow up of patients who received UshStat® in a previous study (TDU13600) Patients will be followed for 15 years after treatment 	<ul style="list-style-type: none"> Primary: IAE Secondary: Change from baseline in ocular safety assessments, Delay in retinal degeneration 	<ul style="list-style-type: none"> SSD: Sep. 2013 DE: 2036

Venglustat (GCS inhibitor) GBA-PD

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
MOVES-PD ACT14820 NCT02906020	Phase 2 Efficacy, safety, pharmacokinetics, and pharmacodynamics of venglustat (GZ402671) in patients with Parkinson's Disease (PD) carrying a glucocerebrosidase gene (GBA) mutation	243	<ul style="list-style-type: none"> Patients with PD carrying a GBA mutation or other prespecified variant. Randomized, Double-blind, Placebo Controlled, Parallel Assignment Part 1: Increasing dose of venglustat administered once per day. Duration: up to 48 weeks outside Japan, and up to 64 weeks in Japan Part 2: venglustat dose determined in Part 1 administered once a day Duration: 5,6-week screening, 52-week Tx period, 104-week follow-up period and 6-week post Tx observation 	<ul style="list-style-type: none"> Primary: Change from baseline in Movement Disorder Society Unified PD Rating Scale Part II and III score Secondary: Change from baseline in PD Cognitive Rating Scale, Movement Disorder Society Unified PD Rating Scale Part I, II, and III score, Hoehn and Yahr score 	<ul style="list-style-type: none"> SSD: Dec. 2016 DE: 2021

Insulin glargine / lixisenatide

Type 2 Diabetes Mellitus (T2DM)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
LIXILAN-G EFC13794 NCT02787551	Phase 3 Efficacy and safety of lixilan vs. GLP-1 receptor agonist in patients with type 2 Diabetes not controlled on GLP-1 RAs + OADs, with an extension period	500	<ul style="list-style-type: none"> Patients with T2DM Randomized, Open-label, Active Controlled, Parallel-group Active comparator: Liraglutide/Exenatide/Exenatide ER/Albiglutide/Dulaglutide, Metformin, pioglitazone and SGLT2 inhibitor if taken prior to the study continued 1st period: up to 2 weeks screening, 26-week Tx period and 3 to 9 days follow-up post Tx Extension period 26-week extension after the 26-week Tx for the lixiLan arm only, 3-day follow-up post extension 	<ul style="list-style-type: none"> Primary: Change from baseline in HbA1c [Time Frame: Baseline to 26 weeks] Secondary: % of participants reaching HbA1c targets, Change from baseline in FPG, in 7-point SMPG, in 2-hour PPG during standardized meal test, in blood glucose excursion during standardized meal test , in body weight, Symptomatic hypoglycemia, Safety, % of patients requiring rescue therapy 	<ul style="list-style-type: none"> SSD: Jul. 2016 DE: 2018

Insulin glargine / lixisenatide

Type 2 Diabetes Mellitus (T2DM) - Japan

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
LIXILAN JP-L EFC14113 NCT02752412	Phase 3 Efficacy and safety of lixilan compared to insulin glargine with Metformin in Japanese patients with T2DM inadequately controlled on Basal Insulin and Oral Antidiabetic Drugs	534	<ul style="list-style-type: none"> Japanese Patients with T2DM Randomized, Open-label, Active Controlled, Parallel-group, 2- Tx arm Active comparator: insulin glargine Background therapy: Metformin will be continued Study duration: approximately 41 weeks: up to 2-week screening, 12-week run-in, 26-week randomized Tx period and 3-day post Tx follow-up 	<ul style="list-style-type: none"> Primary: Change from baseline in HbA1c Secondary: % of patients reaching HbA1c <7% or ≤6.5%, Change from baseline, in 2-hour PpG, in blood glucose excursion during standardized meal test, in 7-point SMPG profiles (each time point and average daily value), in body weight, in FPG, in daily dose of insulin glargine, % of patients reaching HbA1c <7% with no body weight gain/no documented symptomatic hypoglycemia, % of patients requiring a rescue therapy, hypoglycemic events, AE, Measurement from baseline of anti-lixisenatide antibodies and of anti-insulin antibodies from baseline 	<ul style="list-style-type: none"> SSD: Aug. 2016 DE: 2018

Insulin glargine / lixisenatide

Type 2 Diabetes Mellitus (T2DM)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
LIXILAN-D LPS14860 NCT03434119	Phase 3 Efficacy and safety of SOLIQUA100/33™ compared to Lantus™ in ethnically/racially diverse patients with T2DM	1200	<ul style="list-style-type: none"> Adult patients with T2DM not achieving glycemic control (i.e. HbA1c between 7.5% and 10% (inclusive)) on basal insulin and OADs, and who are Hispanics of any race, non-Hispanic black/African Americans or non-Hispanic Asians Randomized, open-label, active-controlled, multi-center Study duration: 29 weeks (2-week screening, 26-week randomized open-label tx period, 3-day post tx follow-up) 	<ul style="list-style-type: none"> Primary: Change from baseline to Week 26 in HbA1c (%) (overall and within each ethnic/racial subgroup evaluated) Secondary: (within each ethnic/racial subgroup evaluated): Patients with HbA1c<7% at week 26; change in 2-hour post-prandial glucose (PPG); 2-hour glucose excursion; change in body weight; change in insulin glargine dose at Week 26. Hypoglycemia events, AE 	<ul style="list-style-type: none"> SSD: Feb. 2018 DE: 2019

Lixisenatide

Type 2 Diabetes Mellitus (T2DM) Pediatrics

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
TDR14311 NCT02803918	Phase 1 PK and PD of lixisenatide in Pediatric Patients With T2DM not adequately controlled with metformin and/or basal insulin	24	<ul style="list-style-type: none"> Pediatric patients (≥ 10 and < 18 years old with documented T2DM insufficiently controlled with metformin and/or basal insulin Randomized, double-blind, placebo-controlled, dose escalation (3 ascending repeated doses) Study duration: up to 10 weeks including 6-week Tx period with dose escalation every 2 weeks 	<ul style="list-style-type: none"> Primary: AEs, TEAEs, number of patients with anti-lixisenatide Ab, Secondary: lixisenatide PK parameters, PD (plasma glucose AUC-0-4,5 hours) 	<ul style="list-style-type: none"> SSD: May 2017 DE: 2020

Lantus – Toujeo

U300 Type 1 Diabetes Mellitus (T1DM) - Children

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
EDITION JUNIOR EFC13957 NCT02735044	<p>Phase 3</p> <p>Efficacy and safety of a new formulation of insulin glargine (U300) and Lantus® injected once daily in children and adolescents Age 6 - 17 years with T1DM with a 6-month safety extension period</p>	450	<ul style="list-style-type: none"> Children: 6 to 17 years old with T1DM Randomized, Open-label, Parallel-group, 2- Tx arm Active comparator: insulin glargine Study duration: approximately 58 weeks: up to 2-week screening, 6-month comparative Tx period , 6-month comparative extension period and 4-week post Tx follow-up 	<ul style="list-style-type: none"> Primary: Change from baseline in HbA1c Secondary: % of patients with HbA1c values of <7.5% and % of patients with FPG of ≤130 mg/dL (7.2 mmol/L) without any episode of severe and/or documented (SMPG <54 mg/dL; 3.0 mmol/L) symptomatic hypoglycemia during the last 3 months of the main 6-month randomized period, Change from baseline in FPG, Change from baseline in 24-hour mean plasma glucose and in variability of 24-hour mean plasma glucose based on 8-point SMPG profiles, % of patients with hypoglycemia, % of patients with hyperglycemia with ketosis, % of patients with AE 	<ul style="list-style-type: none"> SSD: April 2016 DE: 2018

Sotagliflozin (SGLT-1&2 inhibitor) Type 2 Diabetes Mellitus (T2DM)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
SOTA-MONO (301) T2DM EFC14833 NCT02926937	Phase 3 Efficacy and safety of sotagliflozin vs. placebo in patients with T2DM not currently treated with antidiabetic therapy	400	<ul style="list-style-type: none"> Patients (male and female) with T2D, who are treated with diet and exercise only during the 12 weeks prior to screening Randomized, Double-blind, Placebo-controlled, Parallel-group, 3-Tx arm, sota dose 1/200mg, sota dose 2/400mg, placebo Study duration: up to 34-week: up to 2-week screening period, 2-week single-blind placebo run-in, 26-week double-blind Tx period and 4-week post Tx follow-up 	<ul style="list-style-type: none"> Primary: Change from Baseline in HbA1c in comparison of sotagliflozin dose 1 vs. placebo Secondary: Change from baseline in 2-hour PPG following a mixed meal in comparison of sotagliflozin doses 1/2 vs. placebo, FPG in comparison of sotagliflozin dose 1 vs. placebo, Body weight in comparison of sotagliflozin doses 1/2 versus placebo, % of patients with HbA1c <6.5% in comparison of sotagliflozin dose 1 vs. placebo, % of patients with HbA1c <7.0% in comparison of sotagliflozin dose 1 vs. placebo, Change from Baseline in HbA1c in comparison of sotagliflozin dose 2 vs. placebo, Change from baseline in SBP for patients with baseline SBP ≥130 mmHg in comparison of sotagliflozin dose 1 vs. placebo and SBP for all patients in comparison of sotagliflozin doses 1/2 vs. placebo 	<ul style="list-style-type: none"> SSD: Dec. 2016 DE: 2019

Sotagliflozin (SGLT-1&2 inhibitor) Type 2 Diabetes Mellitus (T2DM)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
SOTA-MET (302) T2DM EFC14834 NCT02926950	Phase 3 Efficacy and safety of sotagliflozin added to metformin in patients with T2DM who have inadequate glycemic control on metformin	500	<ul style="list-style-type: none"> Patients with T2DM currently treated with diet and exercise and on metformin at a stable dose ≥ 1500 mg/day for at least 12 weeks Randomized, Double-blind, Placebo-controlled, Parallel-group, 2-Tx arm (placebo – sota 400mg), On top of metformin Study duration: up to 87-week: up to 2-week screening period, 2-week single-blind placebo run-in, 26-week double-blind core Tx period , 53-week double-blind extension period and 4-week post Tx follow-up 	<ul style="list-style-type: none"> Primary: Change from Baseline in HbA1c Secondary: Change from Baseline I in 2-hour PPG following a mixed meal, in FPG, in body weight % of patients with HbA1c $< 6.5\%$ - % patients with HbA1c $< 7.0\%$ Change from Baseline I in systolic blood pressure (SBP) for patients with baseline SBP ≥ 130 mmHg in SBP for all patients. 	<ul style="list-style-type: none"> SSD: Dec. 2016 DE: 2019

Sotagliflozin (SGLT-1&2 inhibitor) Type 2 Diabetes Mellitus (T2DM)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
SOTA-SU (307) T2DM EFC14835 NCT03066830	Phase 3 Efficacy and safety of sotagliflozin added to a sulfonylurea alone or in combination with metformin in patients with Type 2 Diabetes who have inadequate glycemic control on a sulfonylurea alone or with metformin	500	<ul style="list-style-type: none"> Patients with T2DM treated with a sulfonylurea (≥half the maximum recommended dose as per local label or MTD as monotherapy or in combination with metformin (≥1500 mg per day or MTD) each at a stable dose for at least 12 weeks Randomized, Double-blind, Placebo-controlled, Parallel-group, 2-Tx arm (placebo – sota 400mg) On top of sulfonylurea alone or in combination with metformin Study duration: up to 85-week: up to 2-week screening period, 2-week single-blind run-in, 26-week double-blind core Tx period, 53-week double-blind extension period and 2-week post Tx follow-up 	<ul style="list-style-type: none"> Primary: Change from Baseline in HbA1c Secondary: Change from baseline in FPG, in body weight, in Systolic Blood Pressure (SBP) for patients with baseline SBP ≥130 mmHg, in SBP for all patients, % of patients with HbA1c <6.5%, % of patients with HbA1c <7.0% 	<ul style="list-style-type: none"> SSD: Mar. 2017 DE: 2019

Sotagliflozin (SGLT-1&2 inhibitor) Type 2 Diabetes Mellitus (T2DM)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
SOTA-CKD3 (306) T2DM EFC14837 NCT03242252	Phase 3 Evaluate the efficacy and safety of sotagliflozin in patients with T2DM and Moderate Renal Impairment who have inadequate glycemic control	780	<ul style="list-style-type: none"> Patients with T2DM (drug-naïve or on antidiabetic therapy) and documented moderate renal insufficiency defined by an estimated glomerular filtration rate (based on the 4 variable Modification of Diet in Renal Disease equation) of ≥ 30 and < 60 mL/min/1.73 m² (CKD 3A, 3B) Randomized, Double-blind, Placebo-controlled, Parallel-group, 3-Tx arm (placebo – sota 200mg - sota 400mg) Study duration: up to 60-week: up to 2-week screening period, 2-week single-blind run-in, 52-week randomized Tx period and 4-week post Tx follow-up 	<ul style="list-style-type: none"> Primary: Change in HbA1c for sota dose 1 and sota dose 2 Secondary: Change from Baseline in FPG (doses 1/2) in SBP for patients with baseline SBP ≥ 130 mmHg (doses 1/2), in SBP for all patients (doses 1/2) and in body weight (doses 1/2), % change in UACR for patients with UACR > 30 mg/g (doses 1/2), % of patients with HbA1c less than 6.5% (doses 1/2), % of patients with HbA1c less than 7.0% (doses 1/2), % of patients with AE (doses 1/2) 	<ul style="list-style-type: none"> SSD: Sept. 2017 DE: 2019

Sotagliflozin (SGLT-1&2 inhibitor) Type 2 Diabetes Mellitus (T2DM)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
SOTA-CKD4 (306) T2DM EFC15166 NCT03242018	Phase 3 Evaluate the efficacy and safety of sotagliflozin in patients with T2DM and severe renal impairment who have inadequate glycemic control	276	<ul style="list-style-type: none"> Patients with T2DM (drug-naïve or on antidiabetic therapy) and documented severe renal insufficiency - CKD4 - defined by an estimated glomerular filtration rate equation (based on the 4 variable modification of diet in renal disease equation) of ≥ 15 and < 30 mL/min/1.73 m² Randomized, Double-blind, Placebo-controlled, Parallel-group, 3-Tx arm (placebo – sota 200mg - sota 400mg) Study duration: up to 60-week: up to 2-week screening period, 2-week single-blind run-in, 52-week randomized Tx period and 4-week post Tx follow-up 	<ul style="list-style-type: none"> Primary: Change from Baseline in HbA1c comparing sotagliflozin dose 1 vs. placebo in CKD4 patients Secondary: Change from baseline in HbA1c comparing sotagliflozin dose 2 vs. placebo, in FPG (doses 1/2), in SBP at for patients with SBP greater than or equal to 130 mmHg (doses 1/2), in SBP in all patients (doses 1/2), in body weight (doses 1/2), % change in the UACR for patients with a UACR > 30 mg/g at baseline (doses 1/2), % of patients with HbA1c less than 6.5% (doses 1 and 2), % of patients with HbA1c less than 7.0% (doses 1 and 2), N of patients with AE (doses 1/2) 	<ul style="list-style-type: none"> SSD: Sept. 2017 DE: 2019

Sotagliflozin (SGLT-1&2 inhibitor) Type 2 Diabetes Mellitus (T2DM)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
SOTA-INS (312) T2DM EFC14868 NCT03285594	Phase 3 Efficacy and safety of sotagliflozin in patients with T2DM who have inadequate glycemic control on Basal Insulin alone or in addition to Oral Antidiabetes Drugs (OADs)	560	<ul style="list-style-type: none"> Patients with T2DM using any types of basal insulin alone or in combination with up to 2 OADs Randomized, Double-blind, Placebo-controlled, Parallel-group, 3-Tx arm (placebo – sota 200mg - sota 400mg) Background therapy with insulin glargine (Lantus®) (with or without OADs) throughout the study Study duration: up to 60-week: up to 2-week screening period, 4-week Lantus® titration single-blind placebo run-in period, 52-week double-blind Tx period and 2-week post Tx follow-up 	<ul style="list-style-type: none"> Primary: Absolute change in hemoglobin A1c (HbA1c) (for sotagliflozin dose 1) Secondary: Change in FPG (for sotagliflozin doses 1/2), in Body Weight (for sotagliflozin doses 1/2), in HbA1c (for sotagliflozin dose 1/2), in SBP for patients with baseline SBP ≥130 mmHg (for sotagliflozin doses 1/2), in SBP for all patients (for sotagliflozin dose 1), 	<ul style="list-style-type: none"> SDD: Oct. 2017 DE: 2019

Sotagliflozin (SGLT-1&2 inhibitor) Type 2 Diabetes Mellitus (T2DM)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
SCORED (303) T2DM EFC14875 NCT03315143	Phase 3 Effects of sotagliflozin on CV and renal events in patients with T2DM, CV risk factors and moderately impaired renal function	10 500	<ul style="list-style-type: none"> Patients : T2DM with glycosylated hemoglobin (HbA1c) \geq 7%, Estimated glomerular filtration rate (eGFR) \geq 25 and \leq 60 mL/min/1.73 m², Age 18 years or older with at least one major CV risk factor or age 55 years or older with at least two minor CV risk factors Randomized, Double-blind, Placebo-controlled, Parallel-group, 2-Tx arm (placebo - sota 400mg) Total Study duration: approximately 27 to 51 months, 24-month recruitment and 27-month of follow-up after the last patient randomized 	<ul style="list-style-type: none"> Primary: Baseline to approx. 51 months, Time to the first occurrence of any of the following clinical events: CV death, Non-fatal myocardial infarction, Non-fatal stroke, Time to the first occurrence of any of the following clinical events: CV death; Hospitalization for heart failure Secondary: Baseline to approx. 51 months, Time to first composite renal event, Time to first composite renal event in subgroup of patients with macroalbuminuria, Total N of heart failure events, CV death , All cause mortality 	<ul style="list-style-type: none"> SSD: Nov. 2017 DE: 2022

Sotagliflozin (SGLT-1&2 inhibitor) Type 2 Diabetes Mellitus (T2DM)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
GLIM (304) T2DM EFC14838 NCT03332771	Phase 3 Efficacy and safety of sotagliflozin vs. glimepiride and placebo in patients with T2DM that are taking metformin monotherapy	930	<ul style="list-style-type: none"> Patients : Patients with T2DM treated with metformin at a stable dose ≥ 1500 mg/day or MTD (documented) for at least 12 weeks prior to screening visit Randomized, Double-blind, Double-dummy, Active and Placebo-controlled, Parallel-group, 4-Tx arm (placebo – glimepiride, sota dose 1, sota dose 2) Total Study duration: up to 58 weeks including 2-week screening phase, 2-week single-blind placebo run-in, 52-week double-blind Tx period and 2-week post Tx follow-up 	<ul style="list-style-type: none"> Primary: Absolute change in hemoglobin A1c (HbA1c) (for sotagliflozin dose 1) Secondary: Change in Body Weight (for sotagliflozin dose), in HbA1c (for sotagliflozin dose 2), in SBP for patients with baseline SBP ≥ 130 mmHg (for sotagliflozin dose 1), in SBP for all patients (for sotagliflozin dose 1), % of patients with at least one hypoglycemic event (for sotagliflozin dose 1), % of patients with AE 	<ul style="list-style-type: none"> SSD: Nov. 2017 DE: 2019

Sotagliflozin (SGLT-1&2 inhibitor) Type 2 Diabetes Mellitus (T2DM)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
T2DM & Mild to Moderate HTA PDY15010 NCT03462069	Phase 2 Multiple Dose Study to Assess the Intestinal, Metabolic and Cardiovascular Effects of an 8 Weeks Treatment With Sotagliflozin Once a Day (QD) as Compared With Empagliflozin QD in Type 2 Diabetes Mellitus (T2DM) Patients With Mild to Moderate Hypertension	40	<ul style="list-style-type: none"> T2 DM patients with Hypertension grades 1 or 2 diagnosed for at least 1 year A Randomized, Double-blind, Parallel-group, 2-treatment Multiple Dose Study Two arms: Treatment A (test): Sotagliflozin 2 tablets administered once daily with 1 empagliflozin placebo capsule prior to the first meal of the day Treatment B (Reference) Empagliflozin 1 capsule administered once daily with 2 sotagliflozin placebo tablets prior to the first meal of the day 	<ul style="list-style-type: none"> Primary: PD parameters in feces (sodium excretion, SCFA, pH), 24-hour urinary: glucose & sodium excretion, 14 h plasma glucose profile and GLP-1 profile after standardized meals Secondary: change in fasting plasma glucose, ABPM, change in plasma aldosterone, change in carotid-femoral pulse wave velocity, CGM, LVEF, change in left ventricular end-diastolic diameter, change in plasma volume, AEs, PK 	<ul style="list-style-type: none"> SSD: Apr. 2018 DE: 2018

Sotagliflozin (SGLT-1&2 inhibitor) Type 2 Diabetes Mellitus (T2DM)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
SOTA-BONE EFC15294 NCT03386344	Phase 3 Efficacy and Bone safety of sotagliflozin in Patients 55 years or older with T2DM and Inadequate Glycemic Control	360	<ul style="list-style-type: none"> Patients with T2DM managed with diet and exercise only or with a stable antidiabetes regimen for more than 12 weeks, 55 years or older A Randomized, Double-blind, Parallel-group, Three arms: Treatment Sotagliflozin (dose 1 and dose 2), placebo 26-week Tx, with 78-week double blind extension period 	<ul style="list-style-type: none"> Primary: HbA1c, change (dose 1) Secondary: Bone mineral density (BMD) of lumbar spine, total hip, and femoral neck, change (dose 1 and 2), Hb1Ac change (dose 2), BW, FPG, SBP, and % of patients with Hb1Ac < 7%, change (dose 1 and 2), AEs 	<ul style="list-style-type: none"> SSD: Mar. 2018 DE: 2021

Sotagliflozin (SGLT-1&2 inhibitor)

Type 2 Diabetes Mellitus (T2DM)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
SOTA-EMPA EFC14867 NCT03351478	Phase 3 Efficacy and Safety of sotagliflozin vs placebo and empagliflozin in Patients with T2DM who have inadequate glycemic control while taking DPP4 inhibitor alone or with metformin	700	<ul style="list-style-type: none"> Patients with T2DM on DPP4 with or without metformin at a stable dose for at least 12 weeks prior to screening A Randomized, Double-blind, Parallel-group, Three arms: Sotagliflozin, empagliflozin, placebo Study duration: up to 34 weeks, including a screening phase up to 2 weeks, a 2-week run-in phase, a 26-week double-blind Tx period, and a 4-week Tx FU 	<ul style="list-style-type: none"> Primary: HbA1c, change Secondary: SBP in patients with SBP \geq 130mmHg, PPG following mixed meal tolerance test (MMTT), FPG, BW, SBP, patients with Hb1Ac < 6,5%, % of patients with Hb1Ac < 7% 	<ul style="list-style-type: none"> SSD: Dec 2017 DE: 2019

Sotagliflozin (SGLT-1&2 inhibitor) Worsening Heart Failure

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
Worsening Heart Failure PDY15079 NCT03292653	Phase 2 Safety, Tolerability and Pharmacodynamic activity of sotagliflozin in Hemodynamically Stable Patients with Worsening Heart Failure	81	<ul style="list-style-type: none"> Patients: Admitted to the hospital with worsening of heart failure Design: Randomized, double-blind, placebo-controlled study consisting of 3 subsequent cohorts. Cohort 1: sotagliflozin 200 mg (n=10) or placebo (n=5) ; Cohort 2: sotagliflozin 400 mg (n=10) or placebo (n=5): Cohort 3: sotagliflozin 200 mg (n=17), 400 mg (n=17) or placebo (n=17) Treatment duration: In each cohort, study treatment is administered orally for 14 days 	<ul style="list-style-type: none"> Primary: Safety and Tolerability; Pharmacodynamics: Changes in hemoconcentration from baseline to 14 days, Changes in plasma volume from baseline to 14 days Secondary: Change in erythropoietin from baseline to 14 days, Change in NT-proBNP from baseline to 14 days 	<ul style="list-style-type: none"> SSD: Dec. 2017 DE: 2019

Sotagliflozin (SGLT-1&2 inhibitor) Worsening Heart Failure

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
SOLOIST-WHF EFC15156 NCT03521934	Phase 3 Effects of sotagliflozin on Clinical Outcomes in Hemodynamically Stable Patients with T2DM POST Worsening Heart Failure	4000	<ul style="list-style-type: none"> Patients with T2DM, admitted to the hospital or urgent heart failure visit for worsening heart failure Design: Randomized, double-blind, placebo-controlled, parallel - group Two Arms: sotagliflozin, placebo Estimated study duration for a given patient: approximately 3 to 32 months 	<ul style="list-style-type: none"> Primary: Time to 1st occurrence of either CV death or hospitalization for heart failure (HHF) in patients with LVEF < 50%, Time to 1st occurrence of either CV death or HHF in the total patient population Secondary: Total number of CV death, HHF or urgent HF visit (including recurrent events), Time to first occurrence of composite renal endpoint, Time to CV death in patients with LVEF < 50%, Time to CV death in the total patient population, Time to all cause mortality in patients with LVEF < 50%, Time to all cause mortality in the total patient population 	<ul style="list-style-type: none"> SSD: June 2018 DE: 2021

Efpeglenatide (Long acting GLP1-R agonist)

Type 2 Diabetes Mellitus

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
T2 DM EFC14822 NCT03353350	Phase 3 Efficacy and Safety of efpeglenatide in Patients with T2DM Inadequately Controlled with diet and Exercise	400	<ul style="list-style-type: none"> A 56-week, multicenter, double-blind, placebo-controlled, 4 parallel arms, randomized study to demonstrate the superiority of once-weekly injection of efpeglenatide 2, 4, or 6 mg in comparison to placebo in HbA1c change from baseline to Week 30 	<ul style="list-style-type: none"> Primary: change in HbA1c (%) from Baseline to Week 30 Secondary: Change in HbA1c (%) from Baseline to Week 56, Change from Baseline to Weeks 30 and 56 in Fasting plasma glucose and 7-point SMPG profiles, HbA1c <7.0% at Week 30 and Week 56 (Y/N), Change from Baseline to Weeks 30 and 56 in body weight, Safety and immunogenicity 	<ul style="list-style-type: none"> SSD: Dec. 2017 DE: 2020

Efpeglenatide (Long acting GLP1-R agonist)

Type 2 Diabetes Mellitus

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
T2 DM CVOT EFC14828 NCT03496298	Phase 3 Effects of efpeglenatide on Cardiovascular outcomes in high cardiovascular risk T2DM patients	4000	<ul style="list-style-type: none"> T2DM patients with Hb1Ac > 7% with either established cardiovascular disease or renal impairment $25 \leq \text{eGFR} < 60 \text{ mL/min}$ and at least one cardiovascular risk factor Randomized, double-blind, placebo-controlled, parallel-group (efpeglenatide 4mg, 6mg, placebo) Estimated study duration per patient up to 36 months approximately Study is event driven; mean follow up of 2,5 years is expected 	<ul style="list-style-type: none"> Primary: time to first Major Adverse Cardiovascular Event (MACE) Secondary: time to first, Expanded cardiovascular outcome event, Composite renal event, AEs 	<ul style="list-style-type: none"> SSD: Apr. 2018 DE: 2021

SAR341402 (Rapid Acting Insulin) Type 1 Diabetes Mellitus (T1DM)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
PDY15083 NCT03436498	Phase 1 Safety assessment of SAR341402 and NovoLog® used in continuous subcutaneous infusion for Type 1 Diabetes Mellitus Patients	30 (evaluable)	<ul style="list-style-type: none"> Multi-center, randomized, open-label, two-sequence, two-treatment, 2-period, active-controlled, 2 x 4 weeks cross-over study assessing the safety of SAR341402 and NovoLog® used in CSII in patients with Type 1 diabetes mellitus (T1DM). Patients will be randomized 1:1 to sequences of either SAR341402/ NovoLog® or NovoLog®/SAR341402. After completion of the first 4 weeks of treatment, patients on SAR341402 will be switched to NovoLog® and patients on NovoLog® will be switched to SAR341402. The study duration for each patient will be approximately 10 weeks, including a 2-week screening period, 2 treatment periods of 4 weeks each, and 1-day post-treatment safety follow-up period. 	<ul style="list-style-type: none"> Primary: number of patients with infusion set occlusions, defined as infusion set change due to failure to correct hyperglycemia (plasma glucose ≥ 250 mg/dL) by insulin bolus via the insulin pump Secondary: Unexplained hyperglycemia, Intervals for infusion set changes, Number of patients with insulin pump alarms for "non-delivery", Patient observation of infusion set occlusions, AEs and SAEs, Number of patients with hypoglycemic events 	<ul style="list-style-type: none"> SSD: May 2018 DE: 2018

SAR341402 (Rapid Acting Insulin)

Type 1 & 2 Diabetes Mellitus

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
GEMELLI 1 EFC15081 NCT03211858	Phase 3 Comparison of SAR341402 to NovoLog®/NovoRapid® in adult patients with Diabetes also using Insulin Glargine, with a 6-month safety extension period	580	<ul style="list-style-type: none"> Patients with T1DM or T2DM diagnosed for at least 12 months, who have been treated with a multiple daily injection regimen with NovoLog®/NovoRapid® OR insulin lispro (100 U/mL) in the last 6 months prior to screening visit AND insulin glargine (100 U/mL) in the last 6 months prior to screening visit OR insulin detemir (Levemir®) in the last 12 months prior to screening visit Randomized, Open-label, Parallel-group Active comparator: NovoLog®/NovoRapid® Study duration: 54-week per patient: 2-week screening period, 26-week Tx period, 26-week comparative safety extension, 1-day follow-up period 	<ul style="list-style-type: none"> Primary: Change in HbA1c (%) from baseline to Week 26 Secondary: Change in HbA1c, Patients with HbA1c <7%, Change in FPG, Change in mean 24-hour plasma glucose concentration, Change in PPG, Change in 7-point SMPG, Hypoglycemic patients, Hypoglycemic events, Anti-SAR341402/NovoLog/NovoRapid antibody status, Tx-induced, Tx-boosted and Tx-emergent anti-insulin antibodies 	<ul style="list-style-type: none"> SSD: Aug. 2017 DE: 2019

SAR425899 (GLP-1R/GCGR) Type 2 Diabetes Mellitus (T2DM) Overweight/Obese

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
Overweight to Obese patients with T2DM TDR15516 NCT03414736	Phase 1 Safety and tolerability of SAR425899 in overweight to obese patients and T2DM patients not requiring anti-diabetic pharmacotherapy with an optional 6-month safety extension period.	60	<ul style="list-style-type: none"> Overweight and obese patients and T2DM not requiring anti-diabetic pharmacotherapy; HbA1c \leq 7.0%. Randomized, open-label, 3 arm study with SAR425899 (3 different dose escalation regimens) Study duration approximately 12 weeks for main study (up-to 3-week screening period, 8-week treatment period, 3-day follow-up period) and approximately 9 months for those participating in the 6 month safety extension (12 weeks main part and 6 month extension). 	<ul style="list-style-type: none"> Primary: Frequency and severity of GI adverse events (main study and 6 month extension) Secondary: Change in body weight, fasting plasma glucose and HbA1c (main study and 6 month extension). 	<ul style="list-style-type: none"> SSD: Jan. 2018 DE: 2018

SAR425899 (GLP-1R/GCGR) Type 2 Diabetes Mellitus (T2DM) Overweight/Obese

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
Overweight to Obese patients with T2DM PDY15264 NCT03350191	Phase 1 A PET/CT tracer study to investigate SAR425899 binding to the liver and pancreas in overweight to obese patients with T2DM	14	<ul style="list-style-type: none"> Overweight and obese patients with T2DM diagnosed at least 1 year prior to study inclusion Open-label study with treatment duration of 20 days Total study duration approximately 4-7 weeks (including 21 day screening period, 20 day treatment period, and 7-day follow up period). 	<ul style="list-style-type: none"> Primary: % glucagon receptor occupancy in the liver (change of glucagon receptor tracer binding in the liver with SAR425899 between day 1 and day 20). Secondary: % GLP-1 receptor occupancy in the pancreas (Change of GLP-1 receptor tracer binding in the pancreas with SAR425899 between day 1 and day 17) 	<ul style="list-style-type: none"> SSD: Dec. 2017 DE: 2018

SAR425899 (GLP-1R/GCGR)

Overweight to Obese Subjects

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
Overweight to Obese Subjects PDY15012 NCT03376802	Phase 1 Effect of SAR425899 on Energy Expenditure in Obese Subjects	30	<ul style="list-style-type: none"> Randomized, double-blind, placebo-controlled study to assess the effect of repeated doses of SAR425899 on energy expenditure and safety in overweight to obese subjects. Total study duration of 5-8 weeks (including 21 day screening period, 7 day run-in period, 19-day treatment period, and 3-day follow-up period). 	<ul style="list-style-type: none"> Primary: sleep energy expenditure (change from baseline to day 19) Secondary: total daily energy expenditure, resting energy expenditure and basal energy expenditure (change from baseline to day 19). 	<ul style="list-style-type: none"> SSD: Feb. 2018 DE: 2018

SAR425899 (GLP-1R/GCGR) Non-Alcoholic SteatoHepatitis (NASH)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
NASH ACT15067 NCT03437720	Phase 2 Efficacy and Safety of SAR425899 for the treatment of Non-Alcoholic SteatoHepatitis (NASH)	126	<ul style="list-style-type: none"> A multi-center, double-blind, randomized, placebo-controlled study to assess the efficacy and safety of SAR425899 for the treatment of NASH in overweight or obese patients with NASH and with or without T2DM. Total study duration: approximately 64 weeks (including an 8 week screening period, a 52 week treatment period, and a 4 week follow-up period). 	<ul style="list-style-type: none"> Primary: Percentage of patients with resolution of NASH (ballooning component of NAS =0) without worsening of fibrosis score at week 52. Secondary: Percentage of patients who achieve status of no hepatocyte ballooning with lobular inflammation score of 0 or 1 without worsening of fibrosis score at week 52; % of patients who achieve an improvement of fibrosis by at least 1 stage without worsening of the hepatocyte ballooning component of NAS at week 52; change from baseline to week 52 in the overall NAS. 	<ul style="list-style-type: none"> SSD: to be confirmed DE: to be confirmed

Alirocumab (anti-PCSK-9 mAb)

Heterozygous Familial Hypercholesterolemia (HeFH)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
ODYSSEY KIDS DFI14223 NCT02890992	Phase 2 Efficacy and safety of alirocumab in children and adolescents with heFH followed by an extension phase	40	<ul style="list-style-type: none"> Patients with diagnosis of HeFH through genotyping or clinical criteria., 8 to 17 years old, treated with optimal dose of statin +/- other LMT(s) or non-statin LMT(s) if statin intolerant at stable dose for at least 4 weeks prior to screening lipid sampling Open-Label, Sequential, Repeated Dose-Finding Study (6 doses tested) Background therapies: optimal dose of statin with or without other LMT or non-statin LMT if statin intolerant at stable dose Study duration: approximately 16-23 weeks: up to 6 (+1) weeks screening period, 8 weeks open-label Tx period, 6 to 8 weeks follow-up period 	<ul style="list-style-type: none"> Primary: % change in calculated LDL-C Secondary: Absolute change in calculated LDL-C, % change in APO B (Apo B), % change in non-high density LP cholesterol (non HDL-C), % change in Total-C, in LP, in TG, in HDL-C, in Apo A-1, Absolute change in Apo B, in non-HDL-C, in Total C, in Lp(a), in TG, in HDL-C, in Apo A-1, in ratio apo B/Apo A-1, % of participants achieving a calculated LDL-C level lower than 130 mg/dL (3.37 mmol/L), % of participants achieving a calculated LDL-C level lower than 110 mg/dL (2.84 mmol/L) 	<ul style="list-style-type: none"> SSD: Sep. 2016 DE: 2018

Alirocumab (anti-PCSK-9 mAb)

Heterozygous Familial Hypercholesterolemia (HeFH)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
HeFH in Children and Adolescents EFC14643 NCT03510884	Phase 3 Efficacy and safety of alirocumab in children and adolescents with HeFH	150	<ul style="list-style-type: none"> Patients with diagnosis of HeFH, 8 to 17 years old, treated with optimal dose of statin +/- other LMT(s) or non-statin LMT(s) if statin intolerant at stable dose Randomized, double-Blind, placebo-controlled followed by an open label treatment period (2 dose tested) Study duration: approximately 110 weeks (run-in period, if needed,: up to 4 weeks [+2 days], screening period, up to 2 weeks (+5 days), double-blind treatment period: 24 weeks, open label treatment: 80 weeks) 	<ul style="list-style-type: none"> Primary: % change in calculated LDL-C Secondary: % change in LDL-C, % change in APO B (Apo B), % change in non-high density LP cholesterol (non HDL-C), % change in Total-C, patients with LDL-C level lower than 130 mg/dL (3.37 mmol/L), patients with LDL-C level lower than 110 mg/dL (2.84 mmol/L), % change in Lp(a), in HDL-C, in TG and in ApoA-1. Number of AE, maturing cognition (Cogstate battery test) and pubertal development (Tanner stage) 	<ul style="list-style-type: none"> SSD: May 2018 DE: 2022

Alirocumab (anti-PCSK-9 mAb) LDL Lowering China

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
ODYSSEY EAST EFC13889 NCT02715726	Phase 3 Efficacy and safety of alicocumab vs. ezetimibe in Asia in High CV risk patients with Hypercholesterolemia not adequately controlled with their statin therapy	600	<ul style="list-style-type: none"> Patients with hypercholesterolemia and established coronary heart disease (CHD) or CHD risk equivalents who are not adequately controlled with a maximally tolerated daily dose of statin at a stable dose for at least 4 weeks prior to the screening visit (Week -3) Randomized, Double-blind, Parallel Group, 2-Arm Active comparator: ezetimibe Background therapies: atorvastatin, rosuvastatin, or simvastatin continued during the course of the trial Study duration: max 35 weeks: 3-week screening period, 24-week randomized Tx period, 8-week follow-up period 	<ul style="list-style-type: none"> Primary: % change in calculated LDL-C in the intent-to-treat (ITT) population Secondary: % change in calculated LDL-C in the modified ITT (mITT) population, % change in calculated LDL-C, % change in Apo B, in non-HDL-C, in TC, in Lp(a), in HDL-C, in fasting TG, in Apo A-1, % of patients reaching calculated LDL-C <70 mg/dL (1.81 mmol/L) 	<ul style="list-style-type: none"> SSD: Jul. 2016 DE: 2018

Alirocumab (anti-PCSK-9 mAb)

Homozygous Familial Hypercholesterolemia (HoFH)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
ODYSSEY HoFH Regeneron R727-CL-1628 NCT03156621	Phase 3 Evaluate the efficacy and safety of alirocumab in patients with HoFH	54	<ul style="list-style-type: none"> Diagnosis of HoFH by specific genotype or clinical criteria (all patients on LDL apheresis must be diagnosed based on genotype) Randomized, Double-Blind, Placebo-Controlled, Parallel-Group, 2-Arm (alirocumab Q2W, placebo) Study duration: 12-week double-blind Tx period followed by 10-week alirocumab open-label Tx period 	<ul style="list-style-type: none"> Primary: % change in LDL-C ITT population Secondary: % change in Apo B, % change in non-HDL-C, % change in TC, % change in LP(a), % change in HDL-C, % change in fasting TG, % change in Apo A-1, % change in LDL-C, % change in LDL-C, ApoB B, non-HDL-C, TC, Lp(a), HDL-C, fasting TG, Apo A-1 / (m)ITT population, Absolute change in the ratio of Apo B/Apo A-1 (ITT), % of patients with ≥15% reduction in LDL-C, % of patients with ≥30% reduction in LDL-C, % of patients with ≥50% reduction in LDL-C, % of patients with ≥15% reduction, ≥30% reduction, and ≥50% reduction in LDL-C 	<ul style="list-style-type: none"> SSD: Oct. 2017 DE: 2019

Alirocumab (anti-PCSK-9 mAb)

Homozygous Familial Hypercholesterolemia (HoFH)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
HoFH Children and Adolescents EFC14660 NCT03510715	Phase 3 Efficacy and safety of alirocumab in children and adolescents with HoFH	18	<ul style="list-style-type: none"> Patients with diagnosis of HoFH, 8 to 17 years old, treated with optimal dose of statin +/- other LMT(s) or non-statin LMT(s) if statin intolerant at stable dose Single group assignment, open label (2 doses) Study duration: up to 62 weeks, includes (if needed) a run-in period of up to 4 weeks, a screening period of up to 2 weeks, a treatment period of up to 48 weeks, and a follow-up of 8 weeks 	<ul style="list-style-type: none"> Primary: % change in calculated LDL-C Secondary: % change in LDL-C, % change in APO B (Apo B), % change in non-high density LP cholesterol (non HDL-C), % change in Total-C, % change in Lp(a), in HDL-C, in TG and in ApoA-1. Absolute change in LDL-C, number of patients with AE and pubertal development (Tanner stage) 	<ul style="list-style-type: none"> SSD: Aug 2018 DE1: 2020

Alirocumab (anti-PCSK-9 mAb) Neurocognitive Evaluation

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
Neurocognitive Evaluation Regeneron R727-CL-1532 NCT02957682	Phase 4 Evaluate the effect of alirocumab on Neurocognitive function in patients with HeFH and non-HeFH at high and very high cardiovascular risk	2176	<ul style="list-style-type: none"> Patients with hypercholesterolemia and established coronary heart disease (CHD) or CHD risk equivalents who are not adequately controlled with a maximally tolerated daily dose of statin at a stable dose for at least 4 weeks prior to the screening visit Randomized, Double-Blind, Placebo-Controlled, Parallel-Group, 2-Arm (alirocumab Q2W, placebo, 1:1) Study duration: 3 weeks screening, 96-weeks double-blind Tx period 	<ul style="list-style-type: none"> Primary: Change in Cambridge Neuropsychological Test Automated Battery (CANTAB) cognitive domain Spatial Working Memory (SWM) strategy score from baseline to week 96. Secondary (efficacy): % change in calculated LDL-C, % change in Apo B, in non-HDL-C, in TC, in Lp(a), in HDL-C, in fasting TG, in Apo A-1, % of patients reaching calculated LDL-C <70 mg/dL (1.81 mmol/L) and LDL-C < 50mg/dL(1.29 mmol/L). 	<ul style="list-style-type: none"> SSD: Nov 2016 DE: 2020

SAR439152 (Myosin inhibitor)

Obstructive Hypertrophic Cardiomyopathy (oHCM)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
PIONEER-OLE MyoKardia collaboration MYK-461-008 NCT03496168	Phase 2 An Open-Label Extension Study of Mavacamten (MYK-461) in Adults With Symptomatic Obstructive Hypertrophic Cardiomyopathy Previously Enrolled in Study MYK-461-004 (PIONEER)	12	Open label	<ul style="list-style-type: none"> Frequency and severity of adverse events and serious adverse events 	<ul style="list-style-type: none"> SSD: Apr.2018 DE: 2020

SAR439152 (Myosin inhibitor)

Obstructive Hypertrophic Cardiomyopathy (oHCM)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients*	Design	Endpoints*	Status
EXPLORER-HCM MyoKardia Collaboration MYK-461-005 NCT03470545	Phase 3 A Randomized, Double Blind, Placebo Controlled Clinical Study to Evaluate Mavacamten (MYK-461) in Adults With Symptomatic Obstructive Hypertrophic Cardiomyopathy	220	Randomized, double-blind, placebo-controlled	<ul style="list-style-type: none"> Change in exercise capacity (peak VO₂) NYHA functional class 	<ul style="list-style-type: none"> SSD: May 2018 DE: 2020

SAR439152 (Myosin inhibitor)

Non-obstructive Hypertrophic Cardiomyopathy (nHCM)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
MAVERICK-HCM MyoKardia Collaboration MYK-461-006 NCT03442764	Phase 2 A Randomized, Double-blind, Placebo-controlled, Concentration-guided, Exploratory Study of Mavacamten in Patients With Symptomatic Non-Obstructive Hypertrophic Cardiomyopathy (nHCM) and Preserved Left Ventricular Ejection Fraction	60	This is a multicenter, exploratory, randomized, double-blind study of the administration of mavacamten in 60 participants with symptomatic nHCM randomized to receive a 16-week course of mavacamten doses titrated to achieve 1 of 2 target drug concentrations. Dose adjustments will be based on PK parameters.	Primary: Safety and tolerability; Secondary: exercise capacity by peak oxygen uptake (peak VO ₂), changes in NYHA, diastolic and systolic function by echocardiography, symptoms and quality of life measures, NT pro-BNP levels	<ul style="list-style-type: none"> • SSD: March 2018 • DE: 2019

SAR247799 (S1P1 agonist)

Endothelial Function in patients with T2DM

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
Endothelial Function PDY15286 NCT03462017	Phase 1 Study to Assess the Pharmacodynamic Effects of Repeated Oral Doses of SAR247799 on Endothelial Function in Male and Female Patients With Type 2 Diabetes Mellitus	108	<ul style="list-style-type: none"> Type-2 diabetes patients with % flow mediated dilation <7% at screening Treatment groups: SAR247799, placebo, sildenafil (active comparator) Treatment duration: 28 days 	<ul style="list-style-type: none"> Primary: Absolute change from baseline in the % flow-mediated dilation index of brachial artery Secondary: Change from baseline in peak flow induced by acetylcholine iontophoresis measured by laser doppler perfusion monitoring, Safety, PK 	<ul style="list-style-type: none"> SSD: Mar. 2018 DE: 2018

SAR440181 (Myosin activator)

Dilated cardiomyopathy (DCM)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
MYK-491 Phase 1b SAD in DCM Patients MyoKardia Collaboration MYK-491-003 NCT03447990	Phase 1 Randomized, Double-blind, Crossover, Placebo-controlled, Adaptive Design Study of Safety, Tolerability, Preliminary Pharmacokinetics, and Pharmacodynamics of Single Ascending Oral Doses of MYK-491 in Patients With Stable Heart Failure	12	This is a randomized, crossover, double-blind, placebo-controlled, two cohort, sequential ascending single dose study. All patients will receive placebo and active doses of MYK-491 (low, med and/or high).	<ul style="list-style-type: none"> Primary Endpoint: Frequency and severity of treatment-emergent adverse events and serious adverse events. 	<ul style="list-style-type: none"> SSD: Feb. 2018 DE: Nov. 2018

Fitusiran (siRNA targeting Antithrombin/AT3) Hemophilia A & B 1/4

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
Hemophilia A or B LTE14762 ALN- AT3SC-002 NCT02554773	Phase 1/2 Hemophilia A and Hemophilia B Fitusiran in patients with moderate or severe hemophilia A or B	34	<ul style="list-style-type: none"> For patients having participated in a previous fitusiran study Single Group assignment, Open-label 	<ul style="list-style-type: none"> Primary: incidence of treatment-emergent AEs, SAEs, and AEs leading to study drug discontinuation Secondary: Changes in the N of Bleeding Event, the Amount of Factor VIII or Factor IX administered for the Tx of bleeding episodes, health-related QOL plasma levels of antithrombin and thrombin generation 	<ul style="list-style-type: none"> SSD: Sep. 2015 DE: 2021

Fitusiran (siRNA targeting Antithrombin/AT3) Hemophilia A & B 2/4

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
ATLAS-INH EFC14768 ALN- AT3SC-003 NCT03417102	Phase 3 Hemophilia A Hemophilia B Efficacy and Safety of Fitusiran in patients with Hemophilia A or B, with Inhibitory Antibodies to Factor VIII or IX	54	<ul style="list-style-type: none"> In patients suffering from severe hemophilia A or B with inhibitors, Randomized, Parallel Assignment, Open-label Fitusiran and active comparator (on demand bypassing agents) 	<ul style="list-style-type: none"> Primary: Annualized bleeding rate (ABR) [Time Frame: 9 months] Secondary: <ol style="list-style-type: none"> Annualized spontaneous bleeding rate [Time Frame: 9 months] Annualized joint bleeding rate [Time Frame: 9 months] Quality of Life (QOL) as measured by Haem-A-QOL Questionnaire score on a scale of 0-100 with higher scores representing greater impairment. [Time Frame: 9 months] 	<ul style="list-style-type: none"> SSD: Mar. 2018 DE: 2019

Fitusiran (siRNA targeting Antithrombin/AT3) Hemophilia A & B 3/4

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
ATLAS-A/B EFC14769 ALN- AT3SC-004 NCT03417245	Phase 3 Hemophilia A Hemophilia B Efficacy and Safety of Fitusiran in patients with Hemophilia A or B, without Inhibitory Antibodies to Factor VIII or IX	120	<ul style="list-style-type: none"> In patients suffering from severe hemophilia A or B without inhibitors, Randomized, Parallel Assignment, Open-label Fitusiran and active comparator (on demand Factor VIII or IX) 	<ul style="list-style-type: none"> Primary: Annualized bleeding rate (ABR) [Time Frame: 9 months] Secondary: <ol style="list-style-type: none"> Annualized spontaneous bleeding rate [Time Frame: 9 months] Annualized joint bleeding rate [Time Frame: 9 months] Quality of Life (QOL) as measured by Haem-A-QOL Questionnaire score on a scale of 0-100 with higher scores representing greater impairment. [Time Frame: 9 months] 	<ul style="list-style-type: none"> SSD: Jul. 2018 DE: 2019

Fitusiran (siRNA targeting Antithrombin/AT3) Hemophilia A & B 4/4

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
Hemophilia A and B EFC15110 ALN- AT3SC-009 NCT03549871	Phase 3 Fitusiran in patients with severe hemophilia A or B previously receiving bypassing agent prophylaxis	30	<ul style="list-style-type: none"> For patients (child, adult, older adult) with severe hemophilia and inhibitors. Single Group assignment, Open-label 	<ul style="list-style-type: none"> Primary: annualized bleeding rate (ABR) Secondary: annualized spontaneous bleeding rate, annualized joint bleed rate, Quality of Life (QOL) measured by Haem-A-QOL Questionnaire 	<ul style="list-style-type: none"> SSD: May. 2018 (FPI: Q3 2018) DE: 2019

Sutimlimab (BIVV009 - Anti Complement C1s mAb) Cold Agglutinin Disease (CAgD)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
Cardinal BIVV009-03 NCT03347396	Phase 3 Efficacy and Safety of BIVV009 in patients with Primary Cold Agglutinin Disease with a recent history of Blood Transfusion	20	<ul style="list-style-type: none"> Patients suffering from primary cold agglutinin disease (CAD) with at least one blood transfusion within 6 months of enrollment Open-label, Single Group assignment Part A (required for registration): biweekly IV infusion of BIVV009 up to week 26 Part B: long-term safety and durability of response extension phase for patients having completed Part A, BIVV009 dosing for up to 1 year after Part A LPO 	<ul style="list-style-type: none"> Primary (Part A): response rate (no transfusion required and $\geq 2\text{g/dl}$ increase in Hgb); Secondary (Part A): change in bilirubin, change in FACIT-Fatigue Scale Score, change in LDH, number of transfusions and blood units and change in Hgb; Part B: TEAEs, hemoglobin, bilirubin, FACIT-F, LDH, transfusion, haptoglobin, HRU. 	<ul style="list-style-type: none"> SSD: Nov. 2017 DE: 2019 (primary outcome measure)

Sutimlimab (BIVV009 - Anti Complement C1s mAb) Cold Agglutinin Disease (CAgD)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
Cadenza BIVV009-04 NCT03347422	Phase 3 Efficacy and Safety of BIVV009 in patients with Primary Cold Agglutinin Disease without a recent history of Blood Transfusion	40	<ul style="list-style-type: none"> • Patients suffering from primary cold agglutinin disease (CAD) • Randomized, double-blind, placebo controlled • Part A: biweekly IV infusion of BIVV009 or placebo (up to 26 weeks) • Part B: long-term safety and durability of response extension phase for patients having completed Part A. Blinded cross-over loading doses to allow all participants to receive BIVV009 while maintaining Part A blinding. BIVV009 dosing for up to 1 year after Part A L 	<ul style="list-style-type: none"> • Primary (Part A); response rate (no transfusion required and ≥ 1.5 g/dl increase in Hgb); • Secondary (Part A): change in Hgb, change in bilirubin, change in FACIT-Fatigue Scale Score, change in LDH, incidence of symptomatic anemia symptoms • Part B: TEAEs, hemoglobin, bilirubin, FACIT-F, LDH, transfusion, haptoglobin, HRU. 	<ul style="list-style-type: none"> • SSD: Nov. 2017 • DE: 2019 (primary outcome measure)

Sutimlimab (BIVV009 - Anti Complement C1s mAb) Chronic Immune Thrombocytopenia (ITP)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
TNT009-201 NCT03275454	Phase 1 Safety, PK and PD of BIVV009 in patients with Chronic Immune Thrombocytopenia (ITP)	16	<ul style="list-style-type: none"> Patients suffering from chronic ITP with inadequate response to at least 2 prior therapies Open-label, Single Group assignment Part A: Bi-weekly IV infusion of BIVV009 up to 21 weeks Part B: long-term treatment period (for up to one year) for patients who have had benefit from BIVV009 during Part A; patients undergo monitored washout from BIVV009 for up to 9 weeks at end of Part A and enter Part B upon return of thrombocytopenia. 	<ul style="list-style-type: none"> Primary: TEAEs, premature study terminations, Clinical Laboratory Abnormalities Secondary: Change in platelet count; independence from additional ITP therapy; Number of patient who achieve complete response (CR), response (R), and no response (NR); Duration of CR and R; Time to increased platelet count > 30, 50, and 100 x 10⁹/L; number of patients with loss of CR, loss of R: PK parameters, anti-drug antibodies, PD measures (Complement factor measures, thrombopoietin levels, immature platelet fraction, platelet autoantibody/autoantigen) 	<ul style="list-style-type: none"> SSD: Aug. 2017 DE: 2019

BIVV001 (rFVIII Fc-vWF-XTEN*) Hemophilia A

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
EXTEN-A 242HA101 NCT03205163	Phase 1 Phase 2 Safety, Tolerability and PK of a single dose regimen of Single dose of BIVV001 in Previously Treated Adults With Severe Hemophilia A	18	<ul style="list-style-type: none"> Open-Label, Sequential Assignment Low-Dose cohort: low dose of rFVIII** , washout of at least 72-hour and one single low dose of BIVV001 (25 IU/kg) High-Dose cohort: single high dose of rFVIII, washout of at least 96-hour and a single high dose of BIVV001 (65 IU/kg) 	<ul style="list-style-type: none"> Primary: AE's, clinically significant laboratory abnormalities, including inhibitor formation Secondary: PK of rFVIII including Cmax; t½; Vss and AUC∞ 	<ul style="list-style-type: none"> SSD: Jul. 2017 DE: 2019

ST400 (gene-editing technology) Beta-thalassemia

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
Transfusion-dependent Beta-thalassemia (TDT) ST-400-01 NCT03432364	Phase 1/2 Safety, Tolerability and Efficacy of ST400 Autologous Hematopoietic Stem Cell transplant for Tx of Transfusion-Dependent Beta-thalassemia (TDT)	6	<ul style="list-style-type: none"> Patients with clinical diagnosis of TDT with at least 8 documented RBC transfusion events per year and confirmed diagnosis of beta-thalassemia (genetic testing) Open-Label, Single Group Assignment, single dose 	<ul style="list-style-type: none"> Primary: TEAEs Secondary: change from baseline in Hb fractions measurements, change in HbF, change in frequency and volume of packed red blood cells (PRBC) transfusions 	<ul style="list-style-type: none"> SSD: Mar. 2018 DE: (1st) 2020

Caplacizumab - Cablivi™

Acquired Thrombotic Thrombocytopenic Purpura

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
Post-HERCULES ALX0681-C302 NCT02878603	Phase 3 Evaluate the long-term safety and efficacy of caplacizumab, evaluate safety and efficacy of repeated use of caplacizumab and characterize the long-term impact of acquired Thrombotic Thrombocytopenic Purpura (aTTP).	104	<ul style="list-style-type: none"> Prospective follow-up for adult patients (18 years and older) with acquired TTP who completed HERCULES Single group assignment, open label Study duration: Tx period (only for patients who experience a recurrence of aTTP during the study period): initial i.v. dose followed by daily s.c. injections for up to 6 months (max). Total study duration approximately 3 years 	<ul style="list-style-type: none"> Primary: proportion of subjects with TTP-related events, # of and time to TTP-related events, mortality rate, proportion of subjects with, # of and time to recurrence of disease, proportion of subjects with reported major thromboembolic events, # of and time to major thromboembolic events, cognitive function, quality of life assessment, immunogenicity, and AE 	<ul style="list-style-type: none"> SSD: Aug. 2016 DE: 2020

Dengue Vaccine

Co-administration w/ Tdap booster

Asia Pacific Region

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
NCT02992418	Phase 3 Study of a Tetravalent Dengue Vaccine Administered Concomitantly or Sequentially With Adacel® in Healthy Subjects	688	<ul style="list-style-type: none"> Randomized, multicenter, open-label study in 688 subjects aged from 9 to 60 years 	<ul style="list-style-type: none"> Immunogenicity and safety of CYD dengue vaccine and Tdap vaccine when both vaccines are administered concomitantly or sequentially 	<ul style="list-style-type: none"> SSD: Dec. 2016 DE: 2020

Dengue Vaccine

Different schedules

Asia Pacific, Latin America Regions

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
NCT02628444	Phase 2a Immunogenicity and Safety of 3-Dose and Booster Dose of Tetravalent Dengue Vaccine in Healthy Subjects 9 to 50 Years of Age	1050	<ul style="list-style-type: none"> Two-stage, multi-national, multi-center, observer-blind, randomized, placebo-controlled Phase II immunogenicity and safety study of tetravalent dengue vaccine 	<ul style="list-style-type: none"> Immunogenicity and safety of 3-dose primary series and booster dose 	<ul style="list-style-type: none"> SSD: May. 2016 DE: 2020

Dengue Vaccine

Booster dose

Latin America Region

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
NCT02623725	Phase 2b Study of a Booster Dose of a Tetravalent Dengue Vaccine in Subjects Who Previously Completed the 3-dose Schedule	251	<ul style="list-style-type: none"> Multi-center, observer-blind, randomized, placebo-controlled, Phase II trial 	<ul style="list-style-type: none"> Immunogenicity and safety of a booster dose 	<ul style="list-style-type: none"> SSD: Apr. 2016 DE: 2019

Rabies Vaccine

Verorab

Asia Pacific Region

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
NCT01622062	Phase 3 Immunogenicity and Safety of Verorab® in a "One-week" Intradermal Post-exposure Prophylaxis Regimen	600	<ul style="list-style-type: none"> Open-label, randomized, controlled, multi-center, multi-country trial 	<ul style="list-style-type: none"> Immunogenicity and safety of Verorab® in a "One-week" intradermal post-exposure prophylaxis regimen 	<ul style="list-style-type: none"> SSD: Jun. 2012 DE: 2019

Flu Vaccine

Fluzone HD-QIV HV

North America Region

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
NCT03282240	<p>Phase 3</p> <p>Safety and Immunogenicity of High-Dose Quadrivalent Influenza Vaccine in Participants ≥65 Years in the US</p>	2616	<ul style="list-style-type: none"> Ph3 randomized ,modified double blind, active controlled, multi center 	<ul style="list-style-type: none"> Safety, immunogenicity, consistency 	<ul style="list-style-type: none"> SSD: Sep. 2017 DE: 2018

Flu Vaccine

Fluzone HD-QIV HV (Japan)

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
NCT03233217	Phase 1/2 Safety and Immunogenicity of High-Dose Quadrivalent Influenza Vaccine in Patients ≥65 Years	175	<ul style="list-style-type: none"> Ph1/2 randomized, modified double blind, multi center 	<ul style="list-style-type: none"> Safety and immunogenicity 	<ul style="list-style-type: none"> SSD: Sep. 2017 DE: 2018

Meninge Vaccine

MenQuadTT

Greater Europe, Latin America, Asia Pacific Regions

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
NCT03205371	Phase 3 Immunogenicity and Safety of a Meningococcal Conjugate Vaccine Given Concomitantly With Other Vaccines in Toddlers	1200	<ul style="list-style-type: none">Open-label (immunology laboratory technicians will be blinded to group assignment), randomized, parallel-group, active-controlled, multi-center study	<ul style="list-style-type: none">Immunogenicity and safety	<ul style="list-style-type: none">SSD: Nov. 2016DE: 2019

Dengue Vaccine Booster Asia Pacific Region

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
NCT02824198	Phase 2b Immunogenicity and Safety of a Tetravalent Dengue Vaccine Booster Injection in Subjects Who Previously Completed a 3-dose Schedule	260	<ul style="list-style-type: none"> Multi-center, observer-blind, randomized, placebo-controlled, Phase II non-inferiority trial 	<ul style="list-style-type: none"> Immunogenicity and safety of a booster dose 	<ul style="list-style-type: none"> SSD: Jul. 2016 DE: 2019

Rabies Vaccine

Purified Vero Rabies

North America Region

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
NCT03145766	Phase 2 Immunogenicity and Safety of a Purified Vero Rabies Vaccine	320	<ul style="list-style-type: none"> Multicenter, observer-blind, controlled, randomized, Phase II study 	<ul style="list-style-type: none"> Immunogenicity and safety 	<ul style="list-style-type: none"> SSD: Apr. 2017 DE: 2018

Dengue Vaccine

Co-administration w/ HPV

Latin America Region

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
NCT02979535	Phase 3b Immunogenicity and Safety of a Tetravalent Dengue Vaccine Administered Concomitantly or Sequentially With Cervarix®	480	<ul style="list-style-type: none"> Randomized, open-label, multicenter study 	<ul style="list-style-type: none"> Immunogenicity and safety of a Tetravalent Dengue Vaccine administered concomitantly or sequentially with Cervarix® 	<ul style="list-style-type: none"> SSD: Nov. 2016 DE: 2020

Dengue Vaccine

Co-administration w/ HPV

Asia Pacific Region

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
NCT02993757	Phase 3b Immunogenicity and Safety of a Tetravalent Dengue Vaccine Administered Concomitantly or Sequentially With Gardasil®	528	<ul style="list-style-type: none"> Randomized, open-label, multicenter study 	<ul style="list-style-type: none"> Immunogenicity and safety of a Tetravalent Dengue Vaccine administered concomitantly or sequentially with Gardasil® 	<ul style="list-style-type: none"> SSD: Dec. 2016 DE: 2019

Dengue Vaccine Asia Pacific Region

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
NCT01373281	Phase 3 Study of a Novel Tetravalent Dengue Vaccine in Healthy Children Aged 2 to 14 Years in Asia	10275	<ul style="list-style-type: none"> Randomized, double-blind, controlled, multicenter 	<ul style="list-style-type: none"> Efficacy and safety 	<ul style="list-style-type: none"> SSD: Jun. 2011 DE: 2018

Dengue Vaccine

Latin America Region

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
NCT01374516	<p>Phase 3</p> <p>Study of a Novel Tetravalent Dengue Vaccine in Healthy Children and Adolescents Aged 9 to 16 Years in Latin America</p>	20869	<ul style="list-style-type: none"> Randomized, double-blind, controlled, multicenter 	<ul style="list-style-type: none"> Efficacy and safety 	<ul style="list-style-type: none"> SSD: Jun. 2011 DE: 2018

AcP Primary Africa and Middle East Regions

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
NCT02817451	Phase 4 DTaP-IPV-HB-PRP-T Combined Vaccine as a Primary Series and a 2nd Year of Life Booster in HIV-Exposed Infected and Uninfected	100	<ul style="list-style-type: none"> multicenter, open-label, two-arm study 	<ul style="list-style-type: none"> immunogenicity and safety of 3-dose primary series and booster dose 	<ul style="list-style-type: none"> SSD: Jul. 2016 DE: 2020

Adacel+

North America Region

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
NCT02587520	Phase 1 Study of Tetanus Toxoid, Reduced Diphtheria Toxoid, and Acellular Pertussis Vaccine Adsorbed in Healthy Subjects	1350	randomized, modified double-blinded, multi-center, active comparator, dose and formulation ranging, step-down study,	<ul style="list-style-type: none"> Safety and immunogenicity 	<ul style="list-style-type: none"> SSD: Oct, 2015 DE: 2018

Dengue Vaccine Asia Pacific

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
NCT02827162	<p>Exploratory Phase</p> <p>Association of Host Genetics With Vaccine Efficacy and Study of Immune Correlates of Risk From a Tetravalent Dengue Vaccine</p>	364	<ul style="list-style-type: none"> Exploratory, single-center study 	<ul style="list-style-type: none"> Host generic analysis and correlate of protection 	<ul style="list-style-type: none"> SSD: Mar. 2016 DE: 2018

Dengue Vaccine

Latin America, Asia Pacific Regions

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
NCT02948933	<p>Epidemiology Phase</p> <p>Cohort Event Monitoring for Dengvaxia®, CYD-TDV Dengue Vaccine</p>	30 000	<ul style="list-style-type: none"> Observational 	<ul style="list-style-type: none"> Incidence of selected AEs and SAEs, occurrence and frequency of hospitalized dengue disease and SAEs leading to hospitalization or death 	<ul style="list-style-type: none"> SSD: Dec. 2016 DE: 2026

Flu seasonal Vaccine

Asia Pacific Region

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
NCT03344029	Phase 4 Immunogenicity and Safety of the Shenzhen Trivalent Inactivated Influenza Vaccine Versus a Trivalent Influenza Vaccine Comparator in Chinese Subjects 18 to 59 Years	1 600	<ul style="list-style-type: none"> blind-observer, monocenter, randomized, comparative study 	<ul style="list-style-type: none"> Immunogenicity and safety 	<ul style="list-style-type: none"> SSD: Nov. 2017 DE: 2018

Flu Vaccine

Latin America Region

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
NCT03391193	Phase 3 Immunogenicity and Safety of a Multi-Dose Quadrivalent Influenza Vaccine	360	<ul style="list-style-type: none"> Randomized, open-label, active-controlled, multi-center study in Mexico 	<ul style="list-style-type: none"> Immunogenicity and safety 	<ul style="list-style-type: none"> SSD: Dec. 2017 DE: 2019

Flu seasonal Vaccine

North America Region

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
NCT03308825	Phase 4 Safety and Immunogenicity of Fluzone® Quadrivalent and Fluzone® High-Dose, Influenza Vaccines	240	<ul style="list-style-type: none"> multi-center, open-label trial 	<ul style="list-style-type: none"> Safety and immunogenicity 	<ul style="list-style-type: none"> SSD: Sep. 2017 DE: 2018

AcP Primary Vaccine North America Region

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
NCT00855855	Phase 4 Surveillance Program to Determine Product Specific Rates of Invasive Hib Disease	510 000	<ul style="list-style-type: none">Observational	<ul style="list-style-type: none">Surveillance for Hib disease.	<ul style="list-style-type: none">SSD: Feb. 2009DE: 2019

AcP Primary Vaccine North America Region

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
NCT01129362	Phase 4 Rates of Pertussis Disease Among Persons Receiving Pentacel® or Other Pertussis Vaccines	1 538	<ul style="list-style-type: none"> Observational 	<ul style="list-style-type: none"> Occurrence of pertussis disease, as determined by the Wisconsin Division of Public Health (WDPH). 	<ul style="list-style-type: none"> SSD: May 2010 DE: 2018

MenQuad TT Vaccine

North America Region, Latin America Region

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
NCT03077438	Phase 3 Study of an Investigational Quadrivalent Meningococcal Conjugate Vaccine Administered in Children Aged 2 to 9 Years	1 000	<ul style="list-style-type: none"> modified double-blind, randomized, parallel-group, active-controlled, multi-center trial 	<ul style="list-style-type: none"> Safety and immunogenicity 	<ul style="list-style-type: none"> SSD: Feb. 2017 DE: 2018

MenQuad TT Vaccine

Greater Europe Region

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
NCT02955797	Phase 3 Immunogenicity and Safety of an Investigational Quadrivalent Meningococcal Conjugate Vaccine in Toddlers	918	Modified double-blind, randomized, parallel-group, active-controlled, multi-center trial	<ul style="list-style-type: none"> Immunogenicity and safety 	<ul style="list-style-type: none"> SSD: Feb. 2017 DE: 2018

Meninge Vaccine Asia Pacific Region

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
NCT02864927	Phase 4 Postmarketing Surveillance Study for Use of Menactra® in the Republic of Korea	1 200	<ul style="list-style-type: none">Open, Multi-center, observational, active safety surveillance study	<ul style="list-style-type: none">Occurrence of solicited and unsolicited events	<ul style="list-style-type: none">SSD: Jul. 2016DE: 2019

New Pertussis Vaccine

Latin America Region

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
NCT03147898	Phase Epidemiology Observational Study Describing the Immune Profile Induced By Pertussis Vaccines	120	<ul style="list-style-type: none"> Observational, multicenter trial 	<ul style="list-style-type: none"> Immune response to booster dose 	<ul style="list-style-type: none"> SSD: Apr. 2017 DE: 2019

Flu seasonal Vaccine

North America Region

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
NCT01945424	Phase Epidemiology Sanofi Pasteur Quadrivalent Influenza Vaccine (QIV) Pregnancy Registry	1 000	<ul style="list-style-type: none"> Observational 	<ul style="list-style-type: none"> Pregnancy registry 	<ul style="list-style-type: none"> SSD: Nov. 2016 DE: 2024

Japanese Encephalitis Vaccine Asia Pacific Region

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
NCT02933710	Phase 4 Postmarketing Surveillance Study for IMOJEV® in Republic of Korea	632	<ul style="list-style-type: none"> Multi-center, open, observational, active safety surveillance study. 	<ul style="list-style-type: none"> Occurrence of solicited and unsolicited events 	<ul style="list-style-type: none"> SSD: Jul. 2016 DE: 2019

MenQuad TT Vaccine Booster Greater Europe Region

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
NCT03476135	Phase 3 3 Year Follow-up for Antibody Persistence & Booster in subjects previously vaccinated	188	<ul style="list-style-type: none"> Open label, multicenter study to describe immune persistence of the priming dose and immuno and safety of booster dose 	<ul style="list-style-type: none"> immunogenicity and safety 	<ul style="list-style-type: none"> SSD: Feb. 2018 DE: 2019

MenQuad TT Vaccine

Co administration

North America Region

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
NCT03039686	Phase 3 Safety and Immunogenicity for Infants, with co administration with routine pediatric vaccines	2475	Modified double blind study, randomized, parallel groups, active controlled, multicenter	<ul style="list-style-type: none"> Immunogenicity and safety 	<ul style="list-style-type: none"> SSD: Apr. 2018 DE: 2022

MenQuad TT Vaccine

Alternative schedules

Greater Europe Region

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
NCT03547271	Phase 3 Safety and Immunogenicity for alternative schedules in infants	1540	<ul style="list-style-type: none"> Partially modified double blind, randomized, parallel group, active controlled, multi center 	<ul style="list-style-type: none"> immunogenicity and safety 	<ul style="list-style-type: none"> SSD: Nov. 2018 DE: 2022

Yellow Fever Vaccine

Asia Pacific Region

Immuno-inflammation	Diabetes
Oncology	Cardiovascular
Rare Diseases	Rare Blood Disorders
MS, Neuro, Gene therapy	Vaccines

Study	Description	Patients	Design	Endpoints	Status
NCT03541694	Phase 4 Passive Enhanced Safety Surveillance	660	<ul style="list-style-type: none">Multi-center, prospective, passive ESS.	<ul style="list-style-type: none">Safety	<ul style="list-style-type: none">SSD: Apr. 2018DE: 2018